

Strategic engineering for detecting antimicrobial compounds from *Taxus wallichiana* Zucc. (Himalayan yew)

Priyanka Adhikari¹, Vasudha Agnihotri^{2*} and Anita Pandey^{3*} 

¹ Central Ayurveda Research Institute, Bidhannagar- 700091, Kolkata, West Bengal, India

² G.B. Pant National Institute of Himalayan Environment-Himachal Pradesh Regional Centre, Mohal-Kullu, Himachal Pradesh-175126, India

³ Department of Biotechnology, Graphic Era (Deemed to be University), Clement Town, Dehradun-248002, Uttarakhand, India

* Corresponding authors, E-mail: vasudha@gbpihed.nic.in; anitapandey333@gmail.com

Abstract

Taxus wallichiana Zucc. (Himalayan yew) has been well-documented for therapeutically significant active ingredients. Its bark contains pharmaceutically important compounds i.e., taxol and its derivatives which are well known for their anticancer potential. However, *T. wallichiana* has received limited attention for its equally significant antimicrobial properties. Keeping this background in view, *T. wallichiana* was selected for the detailed investigation of, antimicrobial activities and isolation and characterization of secondary metabolites responsible for antimicrobial activity in different plant parts i.e., needle, bark, and stem extracts. In plate-based bioassays, plants exhibited antimicrobial action against the three main categories of microorganisms (fungi, bacteria, and actinobacteria). Based on the preliminary antimicrobial study, methanol and ethyl acetate extracts, were selected for further experiments. The bioautographic technique was used for identification, and the mobile phase was optimized with the help of a selectivity triangle. After continuous column and thin-layer chromatography, fractions were identified as having good antifungal, antibacterial, and antiactinobacterial activity. These fractions were selected for further characterization using techniques like GC-MS/ LC-MS, and FTIR. These analyses support the identification of several fatty acids, including arachidic acid, behenic acid, palmitic acid, and stearic acid; vitamins (nicotinamide); alkaloids (cinchonine, timolol); amino benzamides (procainamide); carbocyclic sugars (myoinositol); and alkane hydrocarbons (hexadecane), which have antimicrobial activity in *T. wallichiana* needles. The information gathered from this study will help modern medicine make new drug discoveries that combine different active ingredients from medicinal plants to treat a wide range of ailments.

Citation: Adhikari P, Agnihotri V, Pandey A. 2024. Strategic engineering for detecting antimicrobial compounds from *Taxus wallichiana* Zucc. (Himalayan yew). *Medicinal Plant Biology* <https://doi.org/10.48130/mpb-0024-0020>

Introduction

Microorganisms are developing multidrug resistance (MDR) for many antibiotic and medicinally important compounds due to improper usage. This is causing a major threat to society, the economy, and public health^[1]. So alternate sources of drugs are being searched which could be suitable for the microbes MDR properties^[2]. There have been documented efforts in research to identify new, powerful natural or synthetic antimicrobial agents to combat MDR^[3,4]. Recently, it was shown that synthetic cationic and hydrophobic peptides with N-terminal labels, which are linked to the human cathelicidin LL-37 peptide, are effective against both Gram-negative and Gram-positive

infections^[5]. Plant-based secondary metabolites are gaining more attention among natural or synthetic antimicrobial agents. Plants contain biologically active metabolites, known as phytochemicals, that can be extracted from various plant parts, including the leaves, barks, seeds, flowers, and roots, and can be subsequently employed as sources of antimicrobial compounds^[6,7]. The extraction of these compounds can be affected by both the solvent selected for extraction as well as by the extraction method followed. Similarly, the selection of mobile phase solvents for separating phytochemicals from the extract is also very important. Therefore, optimization of extraction method conditions and mobile phase combinations is

required to separate and identify these complex plant-based bioactive secondary metabolites from crude extracts of medicinal plants^[8].

Taxus wallichiana (Zucc.) is a significant evergreen tree of medicinal importance, it is being used to extract the medicine taxol from its bark, and because of that it has drawn a lot of attention. Generally, it can be grown in temperate regions of the Indian Himalayan region (IHR) between 1800 and 3300 meters above sea level^[9]. The Unani and Ayurvedic medical systems have both looked into the medicinal applications of this species. *T. wallichiana* crude extracts also include bioactive metabolites, frequently with potential antioxidant effects. *T. wallichiana* is renowned for its numerous ethno-medical applications^[10]. Asthma and bronchial problems are managed with its needle paste^[11]. Additionally, Himalayan tribal people use tea made from the needles, stem, and bark of Himalayan yew to treat colds, coughs, and hypertension. It has been found that lignan compounds from *T. baccata* heartwood exhibit both antibacterial and cytotoxic properties against a panel of oncology cell lines^[12]. *T. wallichiana* plant part extracts have been researched mainly for their potential for different types of cancer treatment and with little focus on their antibacterial activities^[10,13]. The antimicrobial potential of the *Taxus wallichiana* plant's needles, bark, and stem has been reported, and according to qualitative assessments, actinobacteria, fungi, and bacteria (both Gram-positive and Gram-negative) were all

inhibited by the crude extracts of the plant. The initial findings of the quantitative estimations made to the minimum inhibitory concentration also provided support for it^[14–16]. A detailed study is required for the identification of antimicrobial compounds present in *T. wallichiana* plant part extracts.

The goal of the current work is to separate and collect the fractions of *T. wallichiana*- plant parts extracts for screening and detection of antimicrobial compounds and then characterize the antimicrobial compounds for their identification to better understand their biochemical makeup.

Material and Methods

Study site and sample collection

Plant samples (needle, bark, and stem) were collected from the Jageshwar area, District Almora (29°35′–29°39′ N and 79°59′–79°53′ E) of Uttarakhand, India. A collection of plant samples was submitted to the herbarium records of G.B. Pant National Institute of Himalayan Environment, Kosi-Katarmal, Almora, Uttarakhand, India (Voucher number: GBPI 5050), and the plant was identified as an evergreen woody tree of *Taxaceae* family, it was identified by Dr. K. Chandra Sekar working as Scientist "F" in the Institute. The gathered plant samples were ground into a fine powder for the next experiments after being cleaned and allowed to air dry.

Chemicals and Solvents

We acquired HPLC-grade ethanol, methanol, acetone, chloroform, ethyl acetate (E acetate), hexane, and acetic acid from Merck, India. Additionally, we obtained tryptone yeast extract broth, potato dextrose broth, agar, formic acid, benzene, and toluene from Hi-media, India. Furthermore, we purchased arachidic acid, behenic acid, palmitic acid, stearic acid, myoinositol, hexadecane, timolol, nicotinamide, procainamide, and cinchonine from Sigma Aldrich, India. Lastly, the silica gel (mesh 60-120) and TLC plates (Silica gel 60 F254) were procured from Merck Germany."

Microorganisms

Two Gram-positive bacteria, *Bacillus subtilis* (NRRL B30408) and *B. megaterium* (MCC3124).

Two Gram-negative bacteria, *Escherichia coli*, and *Serratia marcescens* (MTCC4822).

Two Actinobacteria, *Nocardia tenirefensis* (MCC2012), and *Streptomyces* sp. (MCC2003).

Five fungi i. e., *Paecilomyces variotii* (ITCC3710), *Aspergillus niger* (ITCC2546), *Fusarium oxysporum* (ITCC4219), *F. solani* (ITCC 5017), and *Trametes hirsuta* (MTCC11397) were used in this study. The microbiology lab of the Institute G.B. Pant National Institute of Himalayan Environment in Almora, Uttarakhand, constructed a microbial culture collection from which these test microorganisms were collected.

Extraction using the maceration method

Two grams of plant material (needle, bark, and stem individually) were combined in a 1:5 ratio (dry powder: solvents) with methanol, ethanol, and ethyl acetate. Using parafilm, the conical flask's mouth was sealed. Using a Remi rotary shaker set to 160 rpm for 48 hours at room temperature, samples were macerated. Each solvent extract was vacuum-dried at 35–40 °C using a vacuum oven (Narang Scientific Works, New Delhi, Model-257) following extraction. The dried extract was then

Detecting antimicrobial compounds from *Taxus wallichiana*

dissolved in 2 mL of the appropriate solvent, kept apart, and used later at 4 °C.

Qualitative and quantitative Phytochemical analysis

Various qualitative tests were performed for phytochemical screening following the method described by Gul et al.^[17] All the plant parts extracts were evaluated for total phenolic content, total flavonoid content, and total tannin content, with slight modification (concerning extract volume) in methods as described by Kumaran and Karunakaran^[18]; Quettier et al.^[19]; and Nwinuka et al.^[20], respectively. The results of total phenolic content are expressed in terms of mg gallic acid equivalent per g dry weight of extract (GAE mg/g dw), flavonoid concentration in terms of mg quercetin equivalent per g dry weight of extract (QE mg/g dw), and tannin content in terms of mg tannic acid equivalent per g dry weight of extract (mg/g dw). The standard equations derived from the calibration curves were utilized for quantification.

Preliminary screening of antimicrobial activity in different plant parts i.e., stem needles and bark

Agar plate-based bioassays employing the disc diffusion method were carried out to estimate the antibacterial potential of *T. wallichiana* plant part extracts qualitatively. Fungal culture suspensions were made in Potato Dextrose (PD) agar, whilst bacterial and actinobacterial culture suspensions were made on Tryptone Yeast extract (TYE) agar. With the use of a glass spreader, 100 µL of each test organism (separately) was evenly distributed on the corresponding agar surface (TYE agar plates for bacteria and actinobacteria, and PD agar plates for fungus). With the use of sterile forceps, sterile 5 mm filter paper (Whatman No. 1) discs were put over the agar surface. The agar disc was covered with 15 µL of extract. After that, the plates were incubated at 25 °C. Based on the zone of inhibition (mm) measurements after 24 h for bacteria and 120 h for actinobacteria and fungus, the findings were recorded. Each experiment was carried out in triplicate.

Determination of antimicrobial compounds in the crude needle extracts

Sequential extraction for TLC bio-autography

Ten grams of powdered dried needles were macerated in a 1:5 (w/v) mixture of hexane, chloroform, ethyl acetate, methanol, and water. Hexane, chloroform, ethyl acetate, methanol, and water were used in increasing order of solvent polarity during the maceration of the same sample. Each solvent extract was vacuum dried after extraction using a vacuum oven (Narang Scientific Works, New-Delhi, Model-257) at 35 to 40 °C. The dried extract was then individually dissolved in 2 mL of each solvent.

Separation and identification of antimicrobial compounds

Column and thin layer chromatography (TLC) extracts with antimicrobial activity were further used to isolate and identify antimicrobial compounds. The selectivity triangle approach, created by Snyder, was used to optimize the mobile phase for the chromatographic apparatus^[21]. Through column chromatography, extracts with antibacterial activity were purified using a glass column (30 2 cm) filled with silica gel (60–120 mesh). Hexane was used to charge the silica bed before a 1 mL sample was put into the column and 10 different mobile phase

Detecting antimicrobial compounds from *Taxus wallichiana*

combinations listed in Table 1 were used to elute the sample. After collecting the eluted fractions, TLC-bioautography was used to further separate the fractions that have antibacterial activity.

Based on the selectivity triangle, a TLC plate (silica gel 60 F254, Merck, Germany) was used to separate the fraction with antibacterial activity using various combinations of mobile phases (Table 2). The TLC plates were examined for antimicrobial activity, and any areas with potential for antimicrobial activity were scraped off and dissolved in ethyl acetate. Finally, utilizing the solvent systems listed in Table 3, column chromatography was performed once more to further purify fractions to isolate molecules responsible for antibacterial activity. Afterward, the above-mentioned procedure for testing the antimicrobial activity was used for testing the antimicrobial properties of the fractions and was evaluated using a qualitative test (the disc diffusion assay) and a quantitative assessment (the minimum inhibitory concentration, or MIC). The same mobile phase from which the fractions were taken was utilized as a control.

Determination of antimicrobial compounds in the selected rich fractions

Gas chromatography and mass spectrometry (GC/MS)

With the use of the Shimadzu GC/MS-QP2010 ultra, certain fractions were examined. The eluted components' electron impact (EI) mass spectrum was captured using a 10 μ L injection volume. The column's (DB35-MS Capillary non-polar column with dimensions of 30 mm \times 0.25 mm ID \times 0.25 μ m film) initial temperature was maintained at 50 $^{\circ}$ C for 2 minutes, after which it was raised to 250 $^{\circ}$ C at a rate of 7 $^{\circ}$ C/min for a hold time of 3 min, and then to 280 $^{\circ}$ C for 18 min at a rate of 10 $^{\circ}$ C/min. The carrier gas (Helium) flow rate was held constant at 1.21 mL/min. The mass range of 50-1000 Da has been scanned. NIST 14 and Willey 8 library spectra tools were used to identify the chemical

components.

Liquid chromatography and mass spectrometry (LC/MS)

Using LC/MS (MS Synaptic GZ HDMS LC-MS UPLC H-Class) with a C18 (1.7 μ m) column, injection volume of 20 μ L, and 0.1% formic acid: acetonitrile: methanol (20:30:50 v/v/v; isocratic mode) as mobile phase, the non-volatile bioactive chemicals included in the extract were identified. The column was kept at a constant temperature of 35 $^{\circ}$ C. The chromatogram and mass spectrum analysis was done and the results were compared to the reference mass spectrum found in the Metlin library and the online tool Massbank.

FTIR (Fourier Transform Infrared Spectroscopy)

The Agilent-Cary 630 FT-IR spectrometer was used to record the FTIR spectra. Using Microlab FTIR software, 20-40 μ L of samples were used for FTIR analysis spanning the frequency range of 650–4000 cm^{-1} at 8 cm^{-1} intervals. Based on the wave number, the spectrum was further examined for the presence of potential functional groups.

Antimicrobial effects of recognized compounds and their detection in needle extract

Following the identification of antimicrobial compounds, all compounds were procured from Sigma, India, and their antimicrobial activity was assessed through the utilization of the technique delineated in an earlier section. Subsequently, the fatty acids in the needle methanolic extracts are measured. Gas chromatography (Chemito GC, Ceres 800 plus) with a Flame Ionization Detector (FID) was used to evaluate the fatty acids. A BP20 column (30 m \times 0.25 mm) in splitless mode was used for the analysis, and 10 μ L of the extract was injected. The stationary phase film thickness was 0.25 mm. The temperatures of the injector and detector were adjusted to 250 $^{\circ}$ C and 290 $^{\circ}$ C, respectively. The flow rate of the carrier gas, helium, was kept constant at 0.75 mL/min. The oven was preheated at 60 $^{\circ}$ C,

Table 1. Solvent strength of mobile phase used for separation of compounds using column chromatography (CC)

Mobile phase I (%)	Solvent strength	Mobile phase II (%)	Solvent strength	Mobile phase III
Hexane (100)	–	E. Acetate (100)	–	0.1 % acetic acid in methanol (5.1)
Hexane (90)	E. acetate (10)	E. acetate (90)	Methanol (10)	
Hexane (80)	E. acetate (20)	E. acetate (80)	Methanol (20)	
Hexane (70)	E. acetate (30)	E. acetate (70)	Methanol (30)	
Hexane (60)	E. acetate (40)	E. acetate (60)	Methanol (40)	
Hexane (50)	E. acetate (50)	E. acetate (50)	Methanol (50)	
Hexane (40)	E. acetate (60)	E. acetate (40)	Methanol (60)	
Hexane (30)	E. acetate (70)	E. acetate (30)	Methanol (70)	
Hexane (20)	E. acetate (80)	E. acetate (20)	Methanol (80)	
Hexane (10)	E. acetate (90)	E. acetate (10)	Methanol (90)	

Table 2. Solvent strength of mobile phase used for separation of compounds using thin layer chromatography (TLC)

Code	Mobile Phase	Ratio	Solvent strength	Separation	Spot	Rf value
M1	Chloroform: E. acetate: formic acid	(5:4:1)	4.4	Yes	2	0.6, 0.7
M2	E. acetate: methanol: Benzene	(2:0.5:2.5)	3.6	Yes	3	0.3, 0.6, 0.68
M3	Ethanol: chloroform	(1:1)	4.2	Yes	4	0.4, 0.56, 0.64, 0.83
M4	Toluene: E. acetate: formic acid	(6:4:0.5)	3.2	Yes	5	0.25, 0.38, 0.43, 0.64 and 0.81
M5	Hexane: acetone: toluene: ethanol	(10:7:7:6)	5.2	No	No	No
M6	Chloroform: acetonitrile	(7:3)	4.6	Yes	3	0.77, 0.8, 0.85
M7	Chloroform: methanol	(7: 1)	3.9	Yes	8	0.67, 0.15, 0.19, 0.32, 0.40, 0.45, 0.62, 0.75
M8	E. acetate: 2-propanol	(95:5)	4.3	No	No	No
M9	Chloroform: E. acetate: methanol	(25:20:5)	4.3	Yes	5	0.25, 0.4, 0.54, 0.68, 0.76
M10	Chloroform: formic acid: acetonitrile	(6:0.5:3.5)	3.3	No	No	No

Table 3. Solvent strength of mobile phase used for separation of compounds present in spot 2 through thin layer chromatography (TLC) and column chromatography (CC)

Code	Mobile Phase	Ratio	Solvent strength
M 11	Hexane: E. acetate	(7:3)	1.32
M 12	Toluene: methanol	(8:2)	2.94
M 13	Methanol: chloroform	(1:1)	4.6
M 14	DMF: DCM: acetonitrile	(5:1:4)	5.83
M 15	Acetic acid: methanol: water	(5:2.5:2.5)	6.77

then raised to 260 °C at a pace of 3 °C per minute, and left there for ten minutes. By comparing the retention durations of the fatty acids to a variety of fatty acid standards, the fatty acids were found.

Using RP-HPLC (Shimadzu LC solution, Japan) and a PDA (Photodiode Array) detector, non-volatile chemicals were examined. Using a C18 reverse-phase column, the active metabolites in methanolic needle extracts were measured. Every sample had an injection volume of 20 µL, and the mobile phase was operated in gradient mode. The following compounds' wavelengths and mobile phase compositions were measured: procainamide (275 nm, methanol: water), nicotinamide (275 nm, 0.1 % TFA in water: acetonitrile: methanol), myoinositol (280 nm, methanol: 0.02 M H₂SO₄), and cinchonine (260 nm, acetonitrile: methanol: phosphoric acid), with flow rate of 1 mL/min for all the compounds, except 1.2 mL/min for nicotinamide, and 0.6 mL/min for myoinositol.

Statistical analysis

Data were collected from all trials in triplicate and were expressed as a mean with standard deviation (SD). GraphPad, Prisma 8 software was used to create the heatmap.

Results and discussion

Phytochemical analysis

The extract yield was higher in the methanolic extract (15 %) followed by ethanol (12 %) and ethyl acetate (9 %). The qualitative analysis of *T. wallichiana* needle, stem, and bark shows (Fig. 1) the presence of phytochemical substances such as phenol, flavanol, flavonoid, tannin, saponin, terpenoid, glycoside, and steroid. Total Phenolic, flavonoids, and tannin content were higher in ethanolic extracts of needles (82.20 ± 0.51 mg GAE/g (dw), 79.72 ± 0.69 mg QRE/g (dw), 12.98 ± 0.34 mg TAE/g (dw)) and methanol (81.37 ± 0.85 mg GAE/g (dw), 54.10 ± 0.58 mg QRE/g (dw), 10.52 ± 0.29 mg TAE/g (dw)) extracts, respectively. In stem, phenolic (52.57 ± 0.39 mg GAE/g (dw)) and tannin (9.09

Detecting antimicrobial compounds from *Taxus wallichiana*

± 0.11 mg TAE/g (dw)) content was higher in methanolic extracts while flavonoids content was higher in ethanolic extracts (38.41 ± 0.68 mg QRE/g (dw)). Likewise for bark, phenolic (68.52 ± 0.72 mg GAE/g (dw)) content, flavonoids (61.57 ± 1.09 mg QRE/g (dw)), and tannin (13.05 ± 0.18 mg TAE/g (dw)) content were higher in ethyl acetate extract. Results are shown in Fig. 2.

Antimicrobial potential of needle, bark, and stem

Using extracts from *T. wallichiana* needle, bark, and stem a preliminary test was conducted to determine which plant part would be most effective for isolation and identification of antimicrobial metabolites. It's interesting to note that all plant part extracts exhibited antimicrobial action against the three types of microorganisms i.e., fungi, actinobacteria, and bacteria. *T. wallichiana* extracts; were found more effective against *Bacillus* species in terms of inhibition, and were less effective against *E. coli*. Between two tested *Bacillus* species, ethanol needle extract demonstrated greater inhibition for *B. subtilis*, whereas needle methanol extract demonstrated greater inhibition for *B. megaterium*. Likewise, needle and bark methanolic extracts showed higher antimicrobial activity for *S. marcescens*, needle and bark ethanolic extracts for *P. chlororaphis*, and needle methanol and ethanol extract for *E. coli*. Methanolic extracts of needles and bark showed significant inhibition against actinobacteria also i.e., *N. tenirefensis* and *Streptomyces* sp., Fig. 3 displays the zone of inhibition (mm) results. *T. wallichiana* parts extracts showed antifungal activity against five fungus species i.e., *P. variotii*, *T. hirsuta*, *F. oxysporum*, *A. niger*, *F. solani*. *P. variotii* showed highest zone of inhibition with needle ethanolic extracts, *T. hirsuta* with bark ethanolic extract, *F. oxysporum*, *A. niger* with needle methanol, and *F. solani* with needle and bark ethyl acetate extracts. Overall, the extent of the zone of inhibition for ethanolic and methanolic plant component extracts demonstrated strong antibacterial activity (Fig. 3).

Screening for the best extracts for further purification studies

Several combinations of mobile phases were tried using the selectivity triangle approach for TLC, column chromatography, and bioautography to separate antimicrobial chemicals in needle fractions. Column chromatography and TLC have been used to identify and further purify a large number of bioactive fractions. The most often used techniques are thin-layer chromatography (TLC) and column chromatography because they are affordable, convenient, and available in a variety of stationary phases^[22].

Based on the results of the preliminary test of phytochemical




Fig. 1 Qualitative phytochemical screening for stem, bark, and needle extracts.


Detecting antimicrobial compounds from *Taxus wallichiana*


Fig. 2 Total phenolic content, Total flavonoids content, and Total tannin content of stem, bark, and needle extracts of *T. wallichiana*.




Fig. 3 Heatmap showing preliminary screening of antimicrobial potential (zone of inhibition (mm)) of needle, stem, and bark of *T. wallichiana*.

and antimicrobial assay, needles were selected for further studies of antimicrobial compound isolation and characterization. There are several bioactive secondary metabolites from various classes included in the plant's crude organic extracts. When extracting desired compounds, it is beneficial to use multiple solvents sequentially, starting from those with low polarity and gradually increasing to high polarity. This approach helps prevent overlapping similar types of secondary metabolites in the desired extracts. Based on that in the current study, we have utilized hexane, chloroform, ethyl acetate, methanol, and water. We found that only the extracts from methanol and ethyl acetate (ME & EAE) displayed antimicrobial activity, while the hexane, chloroform, and aqueous extracts exhibited no such activity.

Based on the results of antimicrobial activity ME & EAE were selected for further fractionation using TLC and column chromatography. It is essential to fully understand the nature of the existing secondary metabolites in the extract before developing the mobile phase for column and thin-layer chromatography. It is essential to fully understand the nature of the substances contained in the extract before developing the mobile phase for column and thin-layer chromatography. In this context, Gas Chromatography-Mass Spectrometry (GC-MS) was conducted for the extracts (ME & EAE) exhibiting antimicrobial activity. The extract's mass spectra revealed the presence of carboxylic sugars, fatty acids, phenols, sterols, and alkanes.

Benzene, propanol, gamma-sitosterol, phenols, and benzoic acid were prevalent in both extracts. The findings mentioned have been previously documented in the essential oil of *Nasturtium officinale*^[23] and in the leaf extract of *Adiantum capillus*^[24].

Mobile phase designing for column chromatography

The secondary metabolites identified in the crude extract were then utilized to create the mobile phases for column chromatography, which was utilized to separate the secondary metabolites further, using a selectivity triangle. In the present study, optimization of the mobile phase was done through a selectivity triangle. It ultimately identifies and quantifies the blend of intermolecular interactions that occur between solutes and solvents/phases^[25]. The basic principle of the selectivity triangle is shown in Fig. 4 and details on the first set of mobile phases are shown in Table 1.

Bioassay-guided fractionation and screening for antimicrobial activity

Ten-bed volumes of mobile phase (details given in Table 1) were passed through the column containing methanol and ethyl acetate extracts. Collected fractions were subjected to testing for antimicrobial activity. The methanol extract fractions exhibited selective antibacterial activity, while the ethyl acetate extract fractions demonstrated robust antimicrobial and antifungal activity. Following the results, the ethyl acetate




Fig. 4 Schematic diagram of selectivity triangle principle for mobile phase optimization

extract fractions were chosen for further bioautography. A schematic diagram explaining the results is shown in Fig. 5.

TLC-bioautography

Isolated fractions with antimicrobial activity were further subjected to separation of antimicrobial compounds using a bioautographic technique.^[26–28] With the help of a selectivity triangle 10 different mobile phases were designed for the separation of compounds through TLC (Table 2). Out of 10 mobile phases, 3 did not show any separation. The mobile phases, which had showed an R_F value between 0.2–0.8 cm, were considered as good (Table 2).

7 mobile phases M1, M2, M3, M4, M6, M7, and M9 (Table 2) were found suitable for the separation of compounds present in the fraction having antimicrobial activity. Based on R_F values, mobile phase M3 and M4 were used for further study. During bio autography out of 4 spots, three spots showed antimicrobial activity. Spot 1, 2, and 3 have antimicrobial activity. Spot 1 and 3 have selective antibacterial activity and spot 2 have antibacterial, antiactinobacterial, and antifungal activity. Therefore, spot 2 was selected for further separations and detection using column chromatography and thin-layer chromatography. Spot 2 was not separated through M3 and M4. Hence, another

set of mobile phases was designed with the help of a selectivity triangle. The solvent strength of the new mobile phase is given in Table 3. The schematic diagram is shown in Fig. 6a.

Screening for the best fractions having potential antimicrobial activity

Ten-bed volumes were passed through the column packed with spot 2. Fractions were collected through 6 different mobile phases, and afterward, all the fraction was then tested for their antimicrobial activity. After this exercise, 3 rich fractions (FA, FB, and FC) were selected based on their antimicrobial activity against all three tested groups including bacteria, actinobacteria, and fungi. The schematic diagram is shown in Fig. 6b and the results on the zone of inhibition are shown in the heatmap of Fig. 7a.

Description of antimicrobial compounds

All of the fractions were subjected to the identification and characterization of antimicrobial compounds after the final three rich fractions (FA, FB, and FC) with antibacterial activity were chosen. Rich fractions (FA, FB, and FC) were subjected to GCMS (for volatile chemicals), LCMS (for non-volatile compounds), and FTIR (for functional group) analyses to characterize antimicrobial substances. A list of identified




Fig. 5 (a) Schematic representation of different subfractions collected from fractionation of methanol and ethyl acetate fraction of *T. wallichiana* needle. (b) Antimicrobial activity of collected subfractions of ethyl acetate extract through column chromatography.


Detecting antimicrobial compounds from *Taxus wallichiana*


Fig. 6 (a) Schematic diagram for finalizing the mobile phase for the TLC. (b) Schematic diagram for the TLC-Bioautography.




Fig. 7 (a) Heatmap showing antimicrobial potential (zone of inhibition (mm)) of final fraction and isolated compounds having antimicrobial activity. PA= palmitic acid, SA= stearic acid, AA= arachidic acid, MY = myoinositol, and HA = hexadecane, PA = procainamide, CN = cinchonine, NA = nicotinamide, TM = timolol. B1) *B. megaterium*, B2) *B. Subtilis*, B3) *E. coli*; and B4) *S. marcescens*. (A and F2) *A. niger*, B and C *P. variotii*, D) *F. oxysporum* E and F) *T. hirsuta*. B) FTIR spectra of rich fraction FA, FB, and FC of *T. wallichiana* needles.

compounds through LCMS and GCMS is given in Table 4. Compounds like benzoic acid, hexadecenoic acid, or palmitic acid were found in all three fractions according to GCMS analysis. Fatty acids, aromatic carboxylic acids, aliphatic hydrocarbons, and sesquiterpenoid alcohols made up the bulk of the compounds that were found. Likewise, the existence of many categories of substances, such as alkaloids, vitamins, flavones, quinones, carboxylic acid, lipids, etc., was revealed by LCMS analysis.

To find out what functional groups were present in the iden-

tified compounds in the purified fraction, FTIR analysis was performed. Based on the fingerprint region (1500–4000 cm⁻¹) and functional groups, FTIR spectra were analyzed. Fig. 7b displays the FTIR spectrum. The functional groups of the compounds identified by GC-MS and LC-MS were compared with the FTIR spectra. All of the fractions share bands between 1000 and 1275 cm⁻¹ (C-O/C-N str) between 3000 and 3100 cm⁻¹ (C-H str in sp² hybridized carbon, such as alkene = C-H str or aromatic C-H str). The bands at 3362 cm⁻¹, which are common between FB and FC, and the bands between 2800 and 3000

Table 4. A list of identified compounds in fractions showed antimicrobial activity

Fraction (FA) Compounds	Fraction (FB) Compounds	Fraction (FC) Compounds
GC-MS		
1,6-Octadine-3-ol	Benzoic acid	Benzenepropanol
Benzoic acid	4-Tetradecene	Megastigmatrienone
Benzene, 1-methoxy 4 - (2 - propenyl)	2,4-Ditert-butylphenol	Ar-tumerone
Phenol, 2-methyl-5-(1-methylethyl)	9-Octadecanoic acid	Mome-inositol
1-Tridecene	Hexadecane	Hexadecanoic acid
Pentadecane	1,4-Dimethyl-2 phenoxybenzene	9-Octadecenoic acid
Succinic acid	E-15-Heptadecenal	Eicosanoic acid
E-14-Hexadecenal	Benzene dicarboxylic acid	Di-n-octyl phthalate
Neophytadiene	Hexadecanoic acid	
Hexadecanoic acid	1-Nonadecene	
1-Nonadecene	Ecosanoic acid	
Docosanoic acid	Octacosanol	
1- Tetradecanol	Phenol, 2,4-bis(1-phenylethyl)	
Benzene dicarboxylic acid	Di-n-octyl phthalate	
LC-MS		
1- aminocyclopropane-1- carbocyclic acid	Nicotinamide	Nicotinamide
Methyl methanethiosulfonate	Cinchonine	Dodecylsulfonilacetic acid
Methionyl-Glycin	Ranitidine	Diphenoxylic acid
Ergothioneine	Psoralenol	Cinchonine
Procainamide	Squamocin B	Ergothioneine
Alcoifosfamide	Trimethaphan	
Timolol	Timolol	
Boviquinone	Trimethaphan	
Lobaric acid	Frangulanine	

cm^{-1} (C-H str in sp^3 hybridized carbon, such as methyl and methylene C-H str), which were common between FA and FB, suggested that N-H stretching may have occurred based on the band's shape, while broadness suggested the possibility of a carboxylic group or ester. Additionally, the tiny hump at 3362 cm^{-1} suggested that FB and FC may include secondary or tertiary amides. In FA and FB, the band about 2850 cm^{-1} suggested the presence of methyl and methylene C-H str; in fraction FC, this was less evident. The only bands found in FC were around 3017 cm^{-1} and the band at 1675.95 cm^{-1} , which are typically caused by the alkenyl C=C stretch/amide and the aromatic C-H stretch, respectively.

Antimicrobial compounds in needle-rich fractions (FA, FB, and FC)

After all the biochemical characterization i.e., GC-MS, LC-MS, and FTIR, antimicrobial compounds were identified in all three fractions (Table 5). Ten compounds i.e., arachidic acid, behenic acid, palmitic acid, and stearic acid (fatty acid); vitamins (nicoti-

namide); alkaloids (cinchonine, timolol); amino benzamides (procainamide); carbocyclic sugars (myoinositol); and alkane hydrocarbons (hexadecane) were identified among all the compounds in the three rich fractions (FA, FB, and FC) that had antibacterial activity. To assess the antibacterial activity of the standard compounds against all the specified microorganisms, plate-based bioassays and minimum inhibitory concentration (MIC) were employed. Results in detail are given in Fig. 7a. Identified fatty acids were found potential antibacterial and antifungal agents. Likewise in some previous reports also using bioassay-guided fractionation, antibacterial fatty acids have been extracted from a variety of plants. Yff et al.^[29] isolated palmitic acid from *Pentania prunelloides* which was effective towards bacteriological contagions. Palmitic and stearic acids are reported in *Labisia pumila* leaves with antibacterial activities^[30,31].

Shafaghath^[32] has reported the antibacterial potential of fatty acids derived from the various parts of the plant *Hypericum*

Table 5. List of compounds identified in the separated fraction (spot 2) having antimicrobial potential

S. No.	Compounds	Formula	Molar Mass	Classification	Concentration in needles mg/g (dw)
GC-FID analysis					
1	Arachidic acid	$\text{C}_{20}\text{H}_{40}\text{O}_2$	312.53 g/mol	Saturated fatty acid	22.94 ± 0.09
2	Behenic acid	$\text{C}_{22}\text{H}_{44}\text{O}_2$	340.58 g/mol	Saturated fatty acid	31.04 ± 0.05
3	Palmitic acid	$\text{C}_{16}\text{H}_{32}\text{O}_2$	256.43 g/mol	Saturated fatty acid	16.81 ± 0.03
4	Stearic acid	$\text{C}_{18}\text{H}_{36}\text{O}_2$	284.48 g/mol	Saturated fatty acid	20.10 ± 0.06
RP-HPLC-PDA analysis					
5	Cinchonine	$\text{C}_{19}\text{H}_{22}\text{N}_2\text{O}$	294.17 g/mol	Alkaloid	3.75 ± 1.21
6	Nicotinamide	$\text{C}_6\text{H}_6\text{N}_2\text{O}$	122.12 g/mol	Vitamin	21.14 ± 0.53
7	Procainamide	$\text{C}_{13}\text{H}_{21}\text{N}_3\text{O}$	235.325 g/mol	aminobenzamides	14.61 ± 0.71
8	Timolol	$\text{C}_{13}\text{H}_{24}\text{N}_4\text{O}_3\text{S}$	316.421 g/mol	Alkaloid	6.31 ± 0.54
9	Myoinositol	$\text{C}_6\text{H}_{12}\text{O}_6$	180.16 g/mol	carbocyclic sugar	1.45 ± 1.01
10	Hexadecane	$\text{C}_{16}\text{H}_{34}$	226.41 g/mol	alkane hydrocarbon	–

Detecting antimicrobial compounds from *Taxus wallichiana*

scabrum (flower, leaf, stem, and seed). Cerdeiras et al.^[33] reported the antibacterial activity of stearic acid, present in the aerial sections of *Ibicella lutea*. In addition to fatty acids, six more compounds i.e., myoinositol, hexadecane, chinchonine, timolol, procainamide, and nicotinamide were present as per MS data and these compounds also showed antibacterial properties along with antifungal activities. These substances have specific antifungal action in addition to strong antibacterial activity. Comparably, reports of myoinositol's antibacterial properties have been made^[34,35]. Hexadecane from *Allium nigrum* has also been shown to have antibacterial activity^[36]. Finally, through this piece of work, a separate fraction of *T. wallichiana* needle extract was obtained with known compounds, which have shown good antimicrobial activity with selected microbial species. This can be further utilized for in-depth studies with other types of microbes. After testing for antimicrobial activity, the compounds in the needle extracts were quantified using GC and HPLC alongside their standard compounds. The results of the quantification are given in Table 5.

Conclusion

Due to the increasing demand for *T. wallichiana* plant parts, there has been a continuous decline in the availability of raw materials sourced from its natural habitats. Consequently, there is a notable scarcity of information concerning various aspects such as the phytochemical composition and antimicrobial properties of *T. wallichiana*. In the current research, a thorough examination was conducted, emphasizing the diversity of phytochemicals, antimicrobial functions, and selection of suitable solvents for the extraction of bioactive components responsible for antimicrobial activity. Needles showed the highest antimicrobial potential among the stem, bark, and needles of *T. wallichiana*. Based on the preliminary data, rich-fractions of needles were subjected to LC-MS, GC-MS, and FTIR analyses, and a total of 10 compounds with antimicrobial potential i.e., myoinositol, hexadecane, cinchonine, procainamide, nicotinamide, palmitic acid, stearic acid, arachidic acid, behenic acid, and timolol have been identified from the needles. It is also suggested that focusing on utilizing the needles, rather than the bark and stem, is optimal for maximizing their antimicrobial potential. Notably, *T. wallichiana*, an evergreen tree, has needles with a lifespan of approximately three and a half months, underscoring the importance of promoting needle usage for sustainable extraction of taxol and other potential secondary metabolites that are biologically active.

Author Contribution

AP and VA conceptualize the work. PA performed the experiments, data interpretation, and paper writing. AP and VA editing and finalization of the manuscript.

Data Availability

The data that support the findings of this study are available on request from the corresponding author

Acknowledgment

The authors are grateful to Director GBP-NIHE Almora, Adhikari et al. *Medicinal Plant Biology* 2024, in press

Uttarakhand, India, for extending facilities. The first author gratefully acknowledges the support from the National Mission on Himalayan Studies (Ministry of Environment, Forest & Climate Change, Govt. of India, New Delhi) for financial support.

Conflict of interest

The authors declare that they have no conflict of interest.

Dates

Received 30 April 2024; Revised 25 July 2024; Accepted 26 July 2024; In press 14 August 2024

References

- Sellera FP, Da Silva, LC, Lincopan N. 2021. Rapid spread of critical priority carbapenemase-producing pathogens in companion animals: a One Health challenge for a post-pandemic world. *Journal of Antimicrobial Chemotherapy* 76:2225–2229
- Pandey, A. and Agnihotri, V. 2015. Antimicrobials from medicinal plants: Research initiatives, challenges, and the future prospects. *Biotechnology of bioactive compounds: sources and applications* 123–150
- Adhikari P, Agnihotri V, Suman SK, Pandey A. 2023. Deciphering the antimicrobial potential of *Taxus wallichiana* Zucc: Identification and Characterization using bioassay guided fractionation. *Chemistry and Biodiversity* 20(1):e202200572
- Eloutify YT, El-Shiekh RA, Ibrahim KM, Elshimy R, et al. 2023. Bioassay-Guided Isolation of Antimicrobial Components and LC/QToF Profile of *Plumeria obtusa*: Potential for the Treatment of Antimicrobial Resistance. *ACS Omega*, 8(7):6476–6491
- Krishnamoorthy R, Adhikari P, Anaikutti P. 2023. Design, synthesis, and characterization of non hemolytic antimicrobial peptides related to human cathelicidin LL-37. *RSC Advances* 13:15594–15605
- Tungmunnithum D, Thongboonyou A, Pholboon A, Yangsabai A. 2018. Flavonoids and other phenolic compounds from medicinal plants for pharmaceutical and medical aspects: an overview. *Medicines* 5(3):93
- Abubakar AR, Haque M. 2020. Preparation of medicinal plants: basic extraction and fractionation procedures for experimental purposes. *Journal of Pharmacy and Bio allied Science* 12(1):1–10
- Zhang Y, Cai P, Cheng G, Zhang Y. 2022. A Brief Review of Phenolic Compounds Identified from Plants: Their Extraction, Analysis, and Biological Activity. *Natural Product Communications* 17(1)
- Nadeem M, Rikhari HC, Kumar A, Palni LMS, Nandi SK. 2002. Taxol content in the bark of Himalayan Yew in relation to tree age and sex. *Phytochemistry* 60:627–631
- Agnihotri, V. , 2023. *Taxus baccata* subsp. *wallichiana* (Zucc.) Pilg. In *Himalayan Fruits and Berries* (pp. 419-434). Academic Press
- Singh H. 2007. 'Himalyan Yew (*Taxus wallichiana* Zucc.) a multi-purpose rare gymnosperm in India. *Indian Forestry* 133:690–696
- Sharma H, Garg MA. 2015. A review of traditional use, phytoconstituents and biological activities of Himalayan Yew *Taxus wallichiana*. *Journal of Integrated Medicine* 13:80–89
- Nisar M, Khan I, Ahmad B, Ali I. et al. 2008. Antifungal and antibacterial activities of *Taxus wallichiana* Zucc. *Journal of Enzyme Inhibition and Medicinal Chemistry* 23(2):256–260
- Juyal D, Thawani V, Thaledi S, Joshi M. 2014. Ethnomedical properties of *Taxus wallichiana* Zucc. (Himalayan Yew). *Journal of Traditional and Complementary Medicine* 4:159–161
- Adhikari P, Pandey A, Agnihotri V, Pandey V. 2018. Selection of solvent and extraction method for determination of antimicrobial potential of *Taxus wallichiana* Zucc. , *Research in Pharmacy* 8:1–9
- Bhardwaj V. 2023. *Taxus wallichiana* Zucc. (Himalayan Yew): A

- Medicinal Plant Exhibiting Antibacterial Properties. *Advances in Experimental Medicine and Biology* 1370:145–153
17. Gul R, Jan SU, Faridullah S, Sherani S, Jahan N. 2017. Preliminary phytochemical screening, quantitative analysis of alkaloids, and antioxidant activity of crude plant extracts from *Ephedra intermedia* indigenous to Balochistan. *The Scientific World Journal* 1:1–7
 18. Kumaran A, Karunakaran RJ. 2007. *In vitro* antioxidant activities of methanol extracts of *Phyllanthus* species from India. *Food Science and Technology* 40(2):344–352
 19. Quettier ACD, Gressier CB, Vasseur AJ, Dine CT. et. el. 2000. Phenolic compounds and antioxidant activities of beak wheat (*Fagopyrum esculentum* Moench) hulls and flour. *Journal of Ethnopharmacology* 72(1-2):35–42
 20. Nwinuka NM, Ibeh GO, Ekeke GI. 2005. Proximate composition and levels of some toxicants in four commonly consumed spices. *Journal of Applied Science and Environmental Management* 9(1):150–155
 21. Snyder LR. 1978. Classification of the solvent properties of the common liquid. *Journal of the Chromatographic Science* 16:223–234
 22. Altemimi A, Lakhssassi N, Baharlouei A, Watson DG, Lightfoot DA. 2017. Phytochemicals: extraction, isolation, and identification of bioactive compounds from plant extracts. *Plants* 6(42):1–23
 23. Mahadevi S, Kheyrollahi M, Sheikhlouei H, Isazzadeh A. 2019. Antibacterial and antioxidant activities of *Nasturtium officinale* essential oil on foodborne bacteria. *The Open Microbiology Journal* 13:81–88
 24. Hussein HM, Hameed IH, Ibraheem OA. 2016. Antimicrobial activity and spectral chemical analysis of methanolic leaves extract of *Adiantum capillus veneris* using GC-MS and FT-IR spectroscopy. *International Journal of Pharmacognosy and Phytochemical Research* 8(3):369–385
 25. Johnson AR, Vitha MF. 2011. Chromatographic selectivity triangle. *Journal of Chromatography A* 1218:556–586
 26. Akhtar MA, Hossain MA, Said SA. 2017. Isolation and characterization of antimicrobial compound from the stem-bark of the traditionally used medicinal plant *Adenium obesum*. *Journal of Traditional and Complementary Medicine* 7:296–300
 27. Habib MR, Karim MR. 2009. Antimicrobial and cytotoxic activity of di-(2-ethylhexyl) phthalate and anhydrosophoradiol- 3-acetate isolated from *Calotropis gigantea* (Linn.) flower. *Mycobiology* 37(1):31–36
 28. Cunico MM, Garcia AC, Lima CP, Cocco LC. 2012. Bioautography to assess antibacterial activity of *Ottonia martiana* Miq. (Piperaceae) on the human oral microbiota. *Journal of Basic and Applied Pharmaceutical Sciences* 33(4):515–519
 29. Yff BTS, Lindsey KL, Taylor MB, Erasmus DG, Jager AK. 2002. The pharmacological screening of *Pentanisia prunelloides* and the isolation of the antibacterial compound palmitic acid. *Journal of Ethnopharmacology* 79:101–107
 30. McGaw LJ, Jager AK, Van-Staden J. 2002. Antibacterial effects of fatty acids and related compounds from plants. *South African Journal of Botany* 68:417–423
 31. Desbois AP, Smith VJ. 2010. Antibacterial free fatty acids: activities, mechanisms of action and biotechnological potential. *Applied Microbiology and Biotechnology* 85(6):1629–1642
 32. Shafaghat A. 2011. Antioxidant, antimicrobial activities and fatty acid components of flower, leaf, stem and seed of *Hypericum scabrum*. *Natural Product Communication* 6:1739–1742
 33. Cerdeiras MP, Fernandez J, Soubes M, Vero S, Ferreira F. et. al. 2000. 'A new antibacterial compound from *Ibicella lutea*. *Journal of Ethnopharmacology* 7:521–525
 34. Brand GD, Magalhaes M, Tinoco M, Aragao F. 2012. Probing protein sequences as sources for encrypted antimicrobial peptides. *PLoS One* 7:1–14
 35. McClean S, Beggs LB, Welch R. 2014. Antimicrobial activity of anti-hypertensive food-derived peptides and selected alanine analogs. *Food Chemistry* 146:443–447
 36. Soussi LSR, Ayeb-Zakhama AE, Mahjoub A, Flamini G. 2014. Chemical composition and antibacterial activity of essential oils from the Tunisian *Allium nigrum* L. *Excli Journal* 13:526–553



Copyright: © 2024 by the author(s). Published by Maximum Academic Press, Fayetteville, GA. This article is an open access article distributed under Creative Commons Attribution License (CC BY 4.0), visit <https://creativecommons.org/licenses/by/4.0/>.