


# Carotenoid cleavage dioxygenase catalyzes carotenoid degradation and regulates carotenoid accumulation and petal coloration in *Zinnia elegans*

Hongsheng Qing<sup>#</sup>, Xinyue Liu<sup>#</sup>, Jiahong Chen, Li Li, Jieyu Qian, Jianxin Fu\* and Chao Zhang\* 

Zhejiang Provincial Key Laboratory of Germplasm Innovation and Utilization for Garden Plants, School of Landscape Architecture, Zhejiang Agriculture and Forestry University, Hangzhou 311300, Zhejiang, China

<sup>#</sup> These authors contributed equally: Hongsheng Qing, Xinyue Liu

\* Corresponding authors, E-mail: [fujx@zafu.edu.cn](mailto:fujx@zafu.edu.cn); [zhangc@zafu.edu.cn](mailto:zhangc@zafu.edu.cn)

## Abstract

Carotenoids are one of the important coloring substances in the ray florets of *Zinnia elegans*. Carotenoid cleavage dioxygenase (CCD) directly affects carotenoid contents by cleaving the double bond of carotenoids, thereby affecting the flower color. However, few reports have elucidated its function in *Zinnia elegans*. In the present study, through transcriptome homology searches, three ZeCCDs, ZeCCD1, ZeCCD4-1, and ZeCCD4-2 were screened from *Zinnia elegans*. The function of ZeCCDs was verified by bacterial pigment complementation system, and it revealed that all of the three ZeCCDs had the ability to cleave  $\beta$ -carotene and other carotenoids ( $\epsilon$ -carotene, zeaxanthin and lycopene). Meanwhile, the expression levels of ZeCCD1 and ZeCCD4-2 in the petals of different cultivars were significantly negatively correlated with the carotenoid content. Furthermore, compared with the cytoplasmic localization of ZeCCD1, ZeCCD4-2 is localized in the plastid. Our results indicate that ZeCCD4-2 is the key gene responsible for the differential accumulation of carotenoids in the petals of different *Zinnia elegans* cultivars. This study lays a foundation for further research on the molecular mechanism of petal coloration and carotenoid metabolism of *Zinnia elegans*.

**Citation:** Qing H, Liu X, Chen J, Li L, Qian J, et al. 2024. Carotenoid cleavage dioxygenase catalyzes carotenoid degradation and regulates carotenoid accumulation and petal coloration in *Zinnia elegans*. *Ornamental Plant Research* 4: e005 <https://doi.org/10.48130/opr-0024-0003>

## Introduction

Carotenoids, a group of widely distributed natural pigments, influence the color development of petals in many plants<sup>[1,2]</sup>. The final accumulation of carotenoids, determined by a series of biosynthetic genes and degradation genes, has a great influence on petal coloration<sup>[3]</sup>. Since the C-C double bond on the carotenoid skeleton is vulnerable to oxygen attack, non-enzymatic or enzymatic oxidative cleavage occurs, resulting in various forms of carotenoid derivatives produced by the enzymatic cleavage of the carotenoid cleavage oxygenases (CCO) family. CCO family composed of multiple members in plants can be divided into two main categories: carotenoid cleavage dioxygenase (CCD) and 9-*cis*-epoxy carotenoid dioxygenase (NCED)<sup>[4]</sup>.

Among different CCD gene members, CCD1 and CCD4 participating in the degradation of carotenoids, are regarded as the key factor to regulate carotenoid accumulation in several plant species<sup>[5-7]</sup>. In plants, the CCD1 enzyme only exists in the cytoplasm due to the lack of plastid localization peptides. Since carotenoids are mainly produced in the plastids, CCD1 cannot directly act on the C<sub>40</sub> carotenoid substrates in the plastids<sup>[8,9]</sup>. Only C<sub>27</sub> apocarotenoids, the product of C<sub>40</sub> carotenoids cleaved by CCD4, can be further cleaved by CCD1 into C<sub>13</sub> and/or C<sub>14</sub> small-molecule volatiles, such as  $\alpha$ -ionone and  $\beta$ -ionone<sup>[10-12]</sup>. But in the bacterial pigment complementation system, when the CCD1 prokaryotic expression vector and the engineering plasmid capable of producing carotenoids are

expressed in *Escherichia coli*, CCD1 can contact and cleave several C<sub>40</sub> carotenoids, including lycopene,  $\alpha$ -carotene,  $\beta$ -carotene,  $\epsilon$ -carotene and zeaxanthin<sup>[13,14]</sup>. Based on these results, it is presumed that CCD1 can directly act on both apocarotenoids produced by other CCD proteins and C<sub>40</sub> carotenoids under certain conditions.

In contrast, the plastid localization of CCD4 allows access to carotenoid substrates that were synthesized and stored in plastids<sup>[15]</sup>. CCD4 cleaves the same position (9, 10 and 9', 10' double bond) of different C<sub>40</sub> carotenoids to produce C<sub>13</sub> apocarotenoids, but CCD4 exhibited different cleavage activity for diversiform C<sub>40</sub> carotenoids, among which  $\beta$ -carotene was considered to be the preferred substrate for CCD4<sup>[16]</sup>. It has been reported that the transcriptional level of CCD4 is a key factor affecting the total carotenoid content, which leads to color differences such as white, pale yellow, and dark orange in plant petals<sup>[17-19]</sup>. For example, the transcription level of *IbCCD4* was negatively correlated with the total carotenoid content during flower coloration changes from yellowish-cream to white in *Lilium brownie* var. *colchesteri*<sup>[20]</sup>. More importantly, in chrysanthemum (*Chrysanthemum morifolium*) petals, there was a significant negative correlation between *CmCCD4a* expression level and total carotenoid content of petals, and white petals turned yellow due to carotenoid accumulation after the RNAi construct of *CmCCD4a* was introduced<sup>[21]</sup>. It is suggested that some carotenoid cleavage dioxygenase genes, rather than carotenoid biosynthesis genes, are largely responsible for carotenoid accumulation in many plant species.

*Zinnia elegans* is an annual herbaceous ornamental flower, which is widely used as summer and autumn flower in domestic and foreign gardens due to its diversity in flower types, color, and versatility of landscaping. Petal color of *Zinnia elegans* ranges widely from red to orange and yellow, which is mainly caused by different content of carotenoids and anthocyanins<sup>[22,23]</sup>.  $\beta$ -carotene is the main carotenoid in the petals, accounting for 65.89% and 82.25% of the total identified carotenoids in red and yellow petals, respectively<sup>[24]</sup>. What's more, the cyclizing function of lycopene cyclase genes intimately associated with  $\beta$ -carotene synthesis, has been verified from *Zinnia elegans*<sup>[24]</sup> by bacterial pigment complementation system<sup>[25]</sup>. However, how the carotenoid cleavage dioxygenase genes, which are closely related to  $\beta$ -carotene degradation, participate in the carotenoid metabolism in the ray florets of *Zinnia elegans* and regulate the molecular mechanism of petal coloration in different color cultivars of *Zinnia elegans* is still unclear and requires further study.

In this study, the red, orange and yellow flower cultivars of *Zinnia elegans*, with different carotenoid content in the petals, were used as study materials. Three CCD genes (one CCD1 and two CCD4) from *Zinnia elegans* were cloned, and further characterized the cleavage ability of these genes on four kinds of C<sub>40</sub> carotenoids using the bacterial pigment complementation system<sup>[25]</sup>. In addition, the carotenoid content and the expression patterns of three CCD genes were also analyzed among the red, orange and yellow cultivars. Finally, the subcellular localization of ZeCCDs was analyzed. This study will help to further explore the molecular mechanism of carotenoid metabolism in *Zinnia elegans*.

## Materials and methods

### Plant materials and growth conditions

The experimental materials were three cultivars from *Zinnia elegans* 'Dreamland' series: *Zinnia elegans* 'Dreamland Red' (DRE), *Zinnia elegans* 'Dreamland Coral' (DC) and *Zinnia elegans* 'Dreamland Yellow' (DY). These plants were cultured in the climate room (photoperiod: 13 h light and 11 h dark, temperature cycle: 23 °C day and 18 °C night) of the College of Landscape Architecture, Zhejiang Agriculture and Forestry University (China). In accordance with previous studies<sup>[22–24]</sup>, the developmental stages of petals were divided into four stages: the elongation stage (S1), the coloring stage (S2), the extension stage (S3) and the blooming stage (S4). The roots, stems, leaves, phyllary and petals of DY at S4 stage were collected for gene expression pattern analysis, and the outermost ray florets at four different developmental stages of DRE, DC and DY were collected to determine total carotenoid content and gene expression patterns. All samples were rapidly collected, frozen in liquid nitrogen, and subsequently stored at –80 °C until analysis.

### Extraction and determination of total carotenoids

Total carotenoid content in petal was determined by spectrophotometric according to Lichtenthaler<sup>[26]</sup>. Fresh petals samples (0.2 g) stored at –80 °C were fully ground, mixed with 95% (v/v) ethanol solution, vortexed, and extracted for 15 min away from light, and centrifuged at 2,500×g for 10 min. The absorbance at 470 nm was measured with the supernatant, and the total carotenoid content of the petals at different

developmental stages of different cultivars were calculated using Lichtenthaler's formula<sup>[26]</sup>.

### Molecular cloning of ZeCCDs

The petal samples of four developmental stages of DY were mixed, and the total RNA in the samples was extracted using UltraClean Polysaccharide and phenol Plant RNA Purification Kit (DNA free) (Novogene, Beijing, China). One  $\mu$ g of high-quality RNA was reverse transcribed into cDNA using the PrimeScript RT Master Mix (Perfect Real Time) Kit (TaKaRa, Dalian, China) for cDNA library construction and sequenced with DNBSEQ platform (Beijing Genomics Institute, Shenzhen, China). CCD1 and CCD4 proteins from *Helianthus annuus* (HaCCD1 XP\_022023037.1; HaCCD4 XP\_021977754.1) was used as bait, and the single gene sequence containing BLASTx (E value < 10<sup>–5</sup>) was searched in the NCBI non-redundant (Nr) protein database ([www.ncbi.nlm.nih.gov](http://www.ncbi.nlm.nih.gov)). Finally, three CCDs (one CCD1 and two CCD4) were obtained, named ZeCCD1, ZeCCD4-1 and ZeCCD4-2, respectively.

The full-length open reading frames (ORFs) of the three ZeCCDs amplified from the petals of DRE, DC, and DY were cloned into the vector pMAL-c5X (New England Biolabs, NEB) using homologous cloning technique (ClonExpress® II One Step Cloning Kit, Vazyme, Nanjing, China) with cloning primers (Supplemental Table S1) to identify gene sequence of each ZeCCD from different cultivars. The recombinant plasmids pMAL-ZeCCDs were transformed into *E. coli* Top10 competent cells and coated on Luria-Bertani (LB) solid medium containing 100  $\mu$ g/mL ampicillin for screening. Single bacteria were selected for Polymerase Chain Reaction (PCR) positive identification and cultured overnight in LB liquid medium containing 100  $\mu$ g/mL ampicillin. The plasmids were extracted and sequenced by Tsingke Biotechnology Co., Ltd (Hangzhou, China).

### In silico analysis

Using the ProtParam tool (<http://web.expasy.org/protparam/>), we analyzed the biochemical properties of the proteins encoded by ZeCCDs from DY, such as protein length, molecular weight (MW), isoelectric point (pI), instability index, aliphatic index and average hydrophilicity (GRAVY). The amino acid sequences of CCDs from *Arabidopsis thaliana*, *Vitis vinifera*, *Triticum aestivum*, *Nicotiana tomentosiformis*, *Zea mays*, *Crocus sativus*, *Crocus angustifolius*, *Crocus chrysanthus*, *Rosa rugosa*, *Osmanthus fragrans*, *Lycium chinense*, *Helianthus annuus*, *Nicotiana tabacum* and *Solanum lycopersicum* were downloaded from GenBank. The GenBank accession numbers of the downloaded sequences are shown in Supplemental Table S2. The sequence alignment was performed with ZeCCDs using MEGA X software<sup>[27]</sup>, and a maximum likelihood phylogenetic tree was constructed based on the JTT matrix-based model (JTT) with 500 bootstrap replicates.

### Functional characterization of ZeCCDs

In order to characterize the cleavage function of ZeCCDs, the above pMAL-ZeCCD1, pMAL-ZeCCD4-1, pMAL-ZeCCD4-2, and the empty vector pMAL-c5X, were respectively co-transformed into *E. coli* Top10 competent cells<sup>[28–30]</sup> with four engineering plasmids (including pAC- $\beta$ ,  $\beta$ -carotene; pAC- $\epsilon$ ,  $\epsilon$ -carotene; pAC-ZEA, zeaxanthin; pAC-LYC, lycopene). The transformed *E. coli* cells were coated on LB solid medium containing 34  $\mu$ g/mL chloramphenicol and 100  $\mu$ g/mL ampicillin, and cultured overnight at 37 °C. Positive single colonies were inoculated in

*ZeCCDs* catalyze carotenoid degradation

2 mL LB liquid medium containing corresponding antibiotics. After 12 h of shaking culture at 37 °C, 1 mL of bacterial culture was expanded to 100 mL of medium containing the same antibiotics, and cultured again at 37 °C for 12 h. The bacterial cells were collected by centrifugation, and carotenoids were extracted with 1 mL of acetone, and shaken until the sediment was colorless. The supernatant was carefully aspirated after centrifugation at 2,500× g for 5 min, filtered through an organic filter membrane with a pore size of 0.22 μm, and stored in a brown glass sample bottle for subsequent carotenoid analysis.

**Expression analysis**

In order to quantitatively analyze the expression of *ZeCCDs* genes, AceQ® qPCR SYBR Green Master Mix (Vazyme, Nanjing, China) was used, and the primers for quantitative real-time PCR (qRT-PCR) are available in Supplemental Table S3. The reaction system and conditions were set according to our previous studies<sup>[24]</sup>. The qRT-PCR analyzes of the target genes were performed on the LightCycler® 480II (Roche Diagnostics, Germany) instrument. With *ZeACT* as the internal reference gene<sup>[22–24]</sup>, Cq values were automatically calculated by LightCycler® 480II software and the relative expression of target genes was quantified by the  $2^{-\Delta Ct}$ <sup>[31]</sup>. All qPCR reactions were set up with three biological replicates.

The total carotenoid content and the expression levels of *ZeCCDs* in the petals of different cultivars at different developmental stages were subjected to Person correlation analysis in SPSS 20.0 software (SPSS Inc., Chicago, IL, USA).

**Subcellular localization analysis**

The coding sequences of *ZeCCDs* from DY without stop codon were inserted into pCNHP-eYFP vector, so that each of the three *ZeCCDs* was fused with the eYFP protein to produce the 35S: *ZeCCDs*-eYFP fusion construct (primers used are shown in Supplemental Table S1). The recombinant plasmids verified by sequencing were transferred into *Agrobacterium tumefaciens* GV3101. The *Agrobacterium tumefaciens* GV3101 containing the target plasmid was respectively injected into 4-week-old tobacco (*Nicotiana benthamiana*) leaves. After 48 h of infection in the dark, the fluorescence signal was observed by confocal fluorescence microscope (ZEISS, Germany) with chloroplast autofluorescence as the marker.


**Results****Carotenoid analysis in petals of different cultivars**

Total carotenoids were extracted from fresh petals of three *Zinnia elegans* cultivars DRE, DC and DY (Fig. 1a) at four developmental stages (Fig. 1b) and quantified by spectrophotometry. The results showed that the total carotenoid content of DY petals was consistently higher than that of DRE and DC at different developmental stages, while that of DRE was always the lowest. Especially at S4, the total carotenoid content in DC petals was 2-fold higher than that of DRE, and the DY content was almost 10-fold higher than that of DRE (Fig. 1c).

**Molecular cloning of *ZeCCDs***

Carotenoid cleavage dioxygenase can breakdown the different sites of different carotenoids, cleave carotenoids, decrease the content of carotenoids in plants, and thus change the plant coloration. Therefore, the isolation, identification and characterization of *ZeCCDs* are essential for understanding carotenoid metabolism in the petals of *Zinnia elegans*. In the current study, the petal transcriptome was used for homology-based search to identify the carotenoid cleavage dioxygenase of *Zinnia elegans*<sup>[24]</sup>. Using *HaCCD1* and *HaCCD4* from *Helianthus annuus* as bait, three homologous genes with the highest sequence identity were identified in *Zinnia elegans*, which were named *ZeCCD1*, *ZeCCD4-1* and *ZeCCD4-2* respectively. Sequence analysis of *ZeCCDs* in DRE, DC and DY showed that sequence identity of *ZeCCD1*, *ZeCCD4-1* and *ZeCCD4-2* in three different cultivars were 100%, 98.21% and 99.78%, respectively. More importantly, we found four histidine residues binding to Fe<sup>2+</sup> cofactors conserved in the amino acid sequences of *ZeCCDs* from three cultivars, which are essential for carotenoid cleavage dioxygenase activity, and their sites in the three cultivars are consistent (Supplemental Fig. S1). Subsequent analysis was carried out with *ZeCCDs* from DY, and coding regions of *ZeCCD1*, *ZeCCD4-1* and *ZeCCD4-2* were 1629 bp, 1734 bp and 1791 bp, respectively (Table 1).

The amino acid sequences of the *ZeCCDs* were compared with protein sequences of *Helianthus annuus*, *Osmanthus fragrans*, *Lycium chinense*, *Arabidopsis thaliana* and other species. It was found that four histidine residues binding to



**Fig. 1** The total carotenoid content in the petals of *Zinnia elegans*. (a) Picture of flowers from DRE, DC and DY, and at (b) different developmental stages. (c) Total carotenoid content in petals from three cultivars and different developmental stages. These are mean ± standard errors, three biological replicates. Different letters indicate significant differences in the one-way ANOVA Duncan test,  $p < 0.05$ .

**Table 1.** Biochemical properties of ZeCCDs.

Protein	ORF length (bp)	Protein length (aa)	Molecular weight (Da)	Isoelectric point	Aliphatic index	Grand average of hydropathy	Instability index
ZeCCD1	1,629	542	61,102.98	5.85	78.54	-0.267	33.71
ZeCCD4-1	1,734	577	64,307.79	6.99	88.01	-0.109	31.13
ZeCCD4-2	1,791	596	65,784.95	6.05	79.13	-0.196	45.14

Fe<sup>2+</sup> cofactors, which are essential for carotenoid cleavage dioxygenase activity, were well conserved in the amino acid sequences of CCDs in the above different specie (Fig. 2a).

Phylogenetic analysis showed that CCDs were mainly divided into two major branches, one of which consisted of CCD1, CCD2 and CCD4. In this branch, CCD1 and CCD4 were clustered into a small branch (Fig. 2b).

### Functional characterization of ZeCCDs

The pAC plasmids, which can provide the genes essential for efficient production of C<sub>40</sub> carotenoids in *E. coli*, were used to establish a system based on the bacterial pigment complementation system<sup>[25]</sup> to verify the cleavage activity of three CCDs on four C<sub>40</sub> carotenoid substrates. Due to the accumulation of carotenoids, cells transformed with different pAC plasmids showed diversiform coloration, which is a convenient system for characterizing enzymes that cleave carotenoids.

The results showed that all three ZeCCDs could make *E. coli* cells, which could produce  $\beta$ -carotene and other carotenoids, show different degrees of discoloration. Compared with pMAL-c5X, the cells co-transformed with four pAC plasmids (pAC- $\beta$ , pAC- $\epsilon$ , pAC-ZEA, pAC-LYC) and pMAL-ZeCCD1 were the lightest in coloration and close to white, while co-transfected with pAC plasmids and pMAL-ZeCCD4 were lighter in coloration (Fig. 3). Carotenoids extracted from *E. coli* cells were examined by spectrophotometer. The results showed that the content of total carotenoids in *E. coli*, which transformed plasmids expressing ZeCCD1 with pAC, was significantly reduced compared to the control, and the content in cells transformed plasmids expressing ZeCCD4-1 and ZeCCD4-2 was also reduced and not significantly different between each other. The above results indicated that these three ZeCCDs in *E. coli* cells showed different cleavage activity for four carotenoids ( $\beta$ -carotene,  $\epsilon$ -carotene, zeaxanthin, lycopene), among which ZeCCD1 had the strongest cleavage activity (Fig. 3).

### Expression patterns of ZeCCDs

The tissue specificity and expression patterns during flower development of ZeCCDs were analyzed by qPCR in order to further understand the possible biological functions of ZeCCDs. In DY, ZeCCD1 had the highest expression level in leaves and the lowest expression level in petals, while ZeCCD4-1 had the highest expression level in petals and stems. The expression level of ZeCCD4-2 was the highest in leaves and phyllaries (Fig. 4a). Significantly, the expression levels of ZeCCD1 and ZeCCD4-2 in petals were much lower than that of ZeCCD4-1.

The results shows that ZeCCD1 has the highest expression level in DRE, followed by DC, and the lowest in DY at all four developmental stages (Fig. 4b). Compared with ZeCCD1, ZeCCD4-1 had the highest expression level in S2, S3 and S4 stages of DY. At the S1 stage, the expression level of ZeCCD4-1 in DC was higher than that in other cultivars (Fig. 4c). The expression pattern of ZeCCD4-2 was similar to that of ZeCCD1. During petal development, both ZeCCD4-2 and ZeCCD1 had the

highest expression level in DRE and the lowest expression level in DY (Fig. 4d).

To further confirm the relationship between the expression levels of ZeCCDs and total carotenoid content, person correlation analysis showed that the expression levels of ZeCCD1 and ZeCCD4-2 were significantly negatively correlated with the total carotenoid content (Table 2).

### Subcellular localization analysis

To determine the subcellular localization of ZeCCDs, we constructed a Pro35S::ZeCCDs-eYFP fusion structure and expressed it in epidermal cells of tobacco leaves, which was then visualized by using fluorescence microscopy. The results of transient expression in tobacco leaves demonstrated that the 35S::ZeCCD1-eYFP fusion protein was localized in the cytosol, while control protein (35S::eYFP) was spread throughout the cell. It is worth noting that the 35S::ZeCCD4-1-eYFP and 35S::ZeCCD4-2-eYFP fusion protein was co-localized with chloroplasts (Fig. 5).

## Discussion


### The amino acid sequences of CCDs were conserved

CCDs can change the type and content of carotenoids in plants by acting on different sites of carotenoids, especially CCD1 and CCD4 which are closely related to plant flower color. In order to explore the metabolic mechanism of carotenoids in the petals of *Zinnia elegans*, three CCDs, including one CCD1 and two CCD4, were identified and isolated from *Zinnia elegans* by using homology-based search on petal transcriptome, and named ZeCCD1, ZeCCD4-1, and ZeCCD4-2 (Table 1). This result was similar to that of *Crocus sativus*, which had two isoforms of CCD4<sup>[32]</sup>. Noteworthy, four histidine residues essential for carotenoid cleavage activity were conserved in these proteins. Four histidine residues together with a divalent iron center form a 4-His-ligated iron center, which is required for substrate recognition and correct positioning of substrates in the active site<sup>[33]</sup>, suggesting that four histidine residues are necessary for CCD to perform its cleavage function. The specific structure of the 4-His-ligated iron center is conserved in the CCD family, which requires the amino acid sequences of CCDs to be evolutionarily conserved<sup>[34]</sup>.


### ZeCCDs can cleave diversiform substrates in *E. coli*

The pigment complementation system in *E. coli* showed that ZeCCD1, ZeCCD4-1 and ZeCCD4-2 cleaved four carotenoids, including  $\beta$ -carotene (Fig. 3), which is the most abundant carotenoids in the petals of *Zinnia elegans*<sup>[24]</sup>. This is similar to the cleavage activity of CCD1 from other plants with extensive affinity for C<sub>40</sub> carotenoids. For example, LeCCD1 from tomato can catalyze the cleavage of several carotenoids such as lycopene,  $\beta$ -carotene,  $\epsilon$ -carotene and zeaxanthin<sup>[10]</sup>. DoCCD1 from *Dendrobium officinale* and RdCCD1 from *Rosa damascena*

ZcCCDs catalyze carotenoid degradation




**Fig. 2** Sequence alignment and phylogenetic analysis of CCDs proteins. (a) Alignment of the deduced amino acid sequences of CCDs homologs. Black represents the 100% homology level of the sequence, dark gray represents the  $\geq 75\%$  homology level, and light gray represents the  $\geq 50\%$  homology level. The black arrow refers to the histidine residue that binds to the  $Fe^{2+}$  cofactor. Sequences of functionally characterized HaCCD1 (XP\_022018126.2) and HaCCD4 (CTP93722.1) from *Helianthus annuus*, OfCCD1 (AXQ60417.1) and OfCCD4 (ABY60887.1) from *Osmanthus fragrans*, LcCCD1 (AIY62758.1) and LcCCD4 (AIY62809.1) from *Lycium chinense*, AtCCD1 (NP\_191911.1) from *Arabidopsis thaliana*. (b) Phylogenetic analysis of CCDs homologs from *Zinnia elegans* and other species (see Supplemental Table S2 for their GenBank accession numbers). The phylogenetic tree was built using MEGA X software with 500 bootstrap replicates based on the JTT matrix-based model (JTT). The scale bar represents the phylogenetic distance calculated based on the number of differences.



**Fig. 3** Functional characterization of ZeCCDs proteins. The pMAL-ZeCCD1, pMAL-ZeCCD4-1, pMAL-ZeCCD4-2 and pMAL-c5X empty vectors were co-transformed into the engineered *E. coli* strains that were modified to accumulate (a)  $\beta$ -carotene, (b)  $\epsilon$ -carotene, (c) zeaxanthin and (d) lycopene. These are mean  $\pm$  standard errors, three biological replicates. Different letters indicate significant differences in the one-way ANOVA Duncan test,  $p < 0.05$ .

also catalyze the cleavage of lycopene,  $\beta$ -carotene and zeaxanthin<sup>[35,36]</sup>. Similarly, two ZeCCD4 have catalytic activity for all four carotenoids, consistent with the reported cleavage activity of CCD4 from *Camellia sinensis* in *E. coli*<sup>[37]</sup>. In contrast, GmCCD4 from soybeans (*Glycine max*) can only specifically catalyze the cleavage of  $\beta$ -carotene and has affinity or cleavage activity for lycopene<sup>[38]</sup>. Although CCD4 enzymes from

different plant species exhibit differences in substrate specificity and regioselectivity, it is generally accepted that  $\beta$ -carotene is recognized as the most favorable substrate<sup>[5]</sup>. Moreover, ZeCCD1 seemed to have higher cleavage activity for diversiform C<sub>40</sub> carotenoids, and only a small quantity of carotenoids could be extracted from *E. coli* co-transformed with different pAC plasmids, compared with ZeCCD4-1 and ZeCCD4-2



**Fig. 4** The expression patterns of ZeCCDs. (a) Expression patterns of three genes in different tissues of DY. The expression patterns of (b) ZeCCD1, (c) ZeCCD4-1 and (d) ZeCCD4-2 in different cultivars and developmental stages. These are mean ± standard errors, three biological replicates. Different letters indicate significant differences between cultivars at the same developmental stage in the one-way ANOVA Duncan test,  $p < 0.05$ .

**Table 2.** Correlation analysis of total carotenoid content and expression levels of ZeCCDs in petals from *Zinnia elegans*.

	ZeCCD1	ZeCCD4-1	ZeCCD4-2
Total carotenoid content	-0.915**	0.543	-0.757*


\* and \*\* indicates significant negative correlation at  $p < 0.05$  and  $p < 0.01$  level between ZeCCDs expression levels and total carotenoid content respectively.

(Fig. 3). This could be partly attributed to the differences of cleavage site and substrate specificity. The CCD1 enzymes are characterized by their extensive substrate and double-bond specificity, and they cleave different linear and cyclic  $C_{40}$  carotenoids and apocarotenoid carotenoids at 9, 10 and 9', 10' positions<sup>[36]</sup>. Accordingly, we surmise that ZeCCD1 catalyzes the cleavage of both ends of carotenoids in two steps, from  $C_{40}$  carotenoids via  $C_{27}$  apocarotenoids to  $C_{13}/C_{14}$  apocarotenoid end-products in *E. coli*, with the second reaction proceeding in a way that may promote the first step. This hypothesis is supported by CCD1 breakdown both  $C_{40}$  carotenoids and  $C_{27}$  apocarotenoids<sup>[11]</sup>. In contrast, ZeCCD4-1 and ZeCCD4-2 may

catalyze single cleavage of  $C_{40}$  carotenoids at 9, 10 (9', 10'), which is similar to research on CCD4 from potato (*Solanum tuberosum*)<sup>[39]</sup>, sweet orange (*Citrus sinensis*)<sup>[40]</sup> and Satsuma mandarin (*Citrus unshiu*)<sup>[41]</sup>. In *Citrus* species, this cleavage led to the production of  $C_{30}$  apocarotenoid pigments, such as  $\beta$ -citraurine and  $\beta$ -citaurin, which are essential for peel pigmentation in citrus fruits<sup>[42]</sup>. Although, it has been reported that CCD4 catalyzes double cleavage of  $C_{40}$  carotenoids at 7, 8 (7', 8'), they are well correlated with the biosynthesis of some specific substances in plants, such as crocins and bixin<sup>[43,44]</sup>, which contain the conjugated polyene system of carotenoids. Therefore, it is necessary to study CCD, which will provide a better understanding of the original biological functions of these enzymes and further determine their hypothetical roles in material metabolism.

### ZeCCD4-2 participates in the cleavage of carotenoids in petals of *Zinnia elegans*

We analyzed the expression patterns of ZeCCDs at four developmental stages in the petals of different *Zinnia elegans* cultivars (Fig. 4). The expression levels of ZeCCD1 and ZeCCD4-2 were significantly negatively correlated with total carotenoid



**Fig. 5** The subcellular localization of ZeCCDs. The fusion constructs (Pro35S::ZeCCDs-eYFP) were transiently transformed into *Nicotiana benthamiana* leaf epidermal cells, and observed by confocal fluorescence microscope (ZEISS, Germany) with chloroplast autofluorescence as the maker. For each construct, yellow (eYFP), red (chloroplasts), gray (bright-field), and merged are shown. Bars = 50 µm.

content in petals of *Zinnia elegans* (Table 2). In general, the expression levels of CCDs are negatively correlated with carotenoid content due to the cleavage function of CCDs on carotenoids. The higher expression level of *SiCCD1* at the grain maturity stage was considered to be the primary reason for the lower carotenoid content in white cultivars of *Setaria italica*<sup>[45]</sup>. What's more, the expression level of *CmCCD4a* in chrysanthemum, which belongs to *Asteraceae* family, also showed a significant negative correlation with the total carotenoid content. After the introduction of an RNAi construct of *CmCCD4a* in the white cultivar, the petals turned yellow<sup>[8]</sup>. The results of the subcellular localization assay showed that ZeCCD1 was localized in the cytosol while ZeCCD4-1 and ZeCCD4-2 were localized in the chloroplasts (Fig. 5). This is consistent with previous studies in *Prunus mume* that PmCCD1 and PmCCD1-like-c were localized in the cytoplasm, and PmCCD4 was localized in the plastid<sup>[46]</sup>. In plants, the enzymes involved in carotenoid biosynthesis are located in the plastids<sup>[35]</sup>. At the same time, carotenoids, as the hypothetical substrate of CCDs, are mostly stored in plastids due to their hydrophobicity, while CCD1 enzyme, due to the lack of plastid localization peptides, is the

only member of the CCDs family located in the cytoplasm. It cannot directly obtain the C<sub>40</sub> carotenoid substrate in the plastid, but acts on the C<sub>27</sub> apocarotenoid, the cleavage product of the C<sub>40</sub> carotenoid that is transported to the cytoplasm<sup>[47]</sup>. Unlike the CCD1 enzymes located in the cytoplasm, most of the CCD4 enzymes are located in the plastids<sup>[13]</sup>, suggesting that the CCD4 enzymes can contact its carotenoid substrate. For example, in Satsuma mandarin (*Citrus unshiu*), CitCCD4 is located in the plastids and can cleave the carotenoid substrates  $\beta$ -cryptoxanthin and zeaxanthin stored in the plastids to produce  $\beta$ -citraurin<sup>[45]</sup>. Therefore, it was presumed that ZeCCD4-2 might be a critical gene affecting carotenoid metabolism in the petals of *Zinnia elegans*. Our work preliminarily revealed the function of carotenoid cleavage dioxygenase genes in carotenoid accumulation in *Zinnia elegans*, and laid a foundation for further exploring the molecular mechanism of carotenoid metabolism in *Zinnia elegans*.

Carotenoid accumulation in plants is not only affected by carotenoid metabolic genes, transcription factors can also be involved in the regulation of carotenoid accumulation. For example, the promoter region of CCD2 in *Crocus sativus*



## ZeCCDs catalyze carotenoid degradation

contained a MYB-binding region, and the expression level of *CstCCD4b* was significantly increased in transgenic lines for the MYB transcription factors *CstMYB1* and *CstMYB1R2*<sup>[48]</sup>. Another study showed that in *Chrysanthemum morifolium*, two MADS box-containing transcription factors, APETALA3 (AP3) and UL1 Interacting Factor 1 (UIF1), could be connected by PISTILLATA (PI) to form the CmAP3-CmPI-CmUIF1 protein complex, regulated carotenoid metabolism by directly activating the expression of *CmCCD4a-2*<sup>[49]</sup>. At the same time, there were many GATA binding sites in the promoter region of *CmCCD4a-5*, and the GATA family transcription factor CmGATA4 of *Chrysanthemum morifolium* could bind directly to the promoter of *CmCCD4a-5* and inhibit its activity<sup>[50]</sup>. The WRKY transcription factor OfWRKY3 and the ethylene responsive factor (ERF) transcription factor OfERF61 of *Osmanthus fragrans* were found to bind directly to the promoter of *OfCCD4* and significantly promote the expression of *OfCCD4*<sup>[51]</sup>. Transcription factors related to the regulation of CCDs in *Zinnia elegans* have not been determined, so more studies should be conducted to identify and reveal the function of up-stream potential transcription factors, which could directly regulate the expression of CCDs in *Zinnia elegans*.

## Conclusions

Three CCDs were isolated from *Zinnia elegans*. These three ZeCCDs could cleave  $\beta$ -carotene,  $\epsilon$ -carotene, zeaxanthin and lycopene. The expression levels of ZeCCD1 and ZeCCD4-2 were significantly negatively correlated with the total carotenoid content in petals from different cultivars. Subcellular localization analysis showed that ZeCCD1 was localized in the cytosol, while ZeCCD4-1 and ZeCCD4-2 were localized in the chloroplasts. Based on the above results, it is speculated that ZeCCD4-2 plays an more important role in carotenoid cleavage in *Zinnia elegans* and affects the petal coloration of ray florets.

## Author contributions

The authors confirm contribution to the paper as follows: literature collation: Qing H; experimental design and implementation: Qing H, Fu J, Zhang C; data analysis and visualization: Qing H, Liu X, Chen J, Li L, Qian J; writing and polishing: Qing H, Liu X; resources and project administration: Fu J, Zhang C; review and editing: Fu J, Zhang C. All authors reviewed the results and approved the final version of the manuscript.

## Data availability

The datasets generated during and/or analyzed during the current study are available from the corresponding author on reasonable request.

## Acknowledgments

We are grateful to Drs E. Gantt and F.X. Cunningham for the *E. coli* functional complementation system and Dr. Xing-Qi Huang for the gift of plasmids pAC-LYC, pAC- $\beta$ , pAC- $\epsilon$ , pAC-Zea and pMAL-C5X. This research was supported by Zhejiang Provincial Natural Science Foundation of China (Grant No. LY24C150001) and College Students Research Training Program of Zhejiang Agriculture and Forestry University (No. 2023KX044).

## Conflict of interest

The authors declare that they have no conflict of interest.

**Supplementary Information** accompanies this paper at (<https://www.maxapress.com/article/doi/10.48130/opr-0024-0003>)

## Dates

Received 26 September 2023; Accepted 5 January 2024; Published online 27 February 2024

## References

- Ohmiya A. 2011. Diversity of carotenoid composition in flower petals. *Japan Agricultural Research Quarterly* 45:163–71
- Zhu C, Bai C, Sanahuja G, Yuan D, Farré G, et al. 2010. The regulation of carotenoid pigmentation in flowers. *Archives of Biochemistry and Biophysics* 504:132–41
- Ohmiya A. 2013. Qualitative and quantitative control of carotenoid accumulation in flower petals. *Scientia Horticulturae* 163:10–9
- Ohmiya A. 2009. Carotenoid cleavage dioxygenases and their apocarotenoid products in plants. *Plant Biotechnology* 26:351–58
- Zheng X, Yang Y, Al-Babili S. 2021. Exploring the diversity and regulation of apocarotenoid metabolic pathways in plants. *Frontiers in Plant Science* 12:787049
- Wang Y, Zhang C, Dong B, Fu J, Hu S, et al. 2018. Carotenoid accumulation and its contribution to flower coloration of *Osmanthus fragrans*. *Frontiers in Plant Science* 9:1499
- Li X, Tang D, Du H, Shi Y. 2018. Transcriptome sequencing and biochemical analysis of perianths and coronas reveal flower color formation in *Narcissus pseudonarcissus*. *International Journal of Molecular Sciences* 19:4006
- Auldrige ME, Block A, Vogel JT, Dabney-Smith C, Mila I, et al. 2006. Characterization of three members of the Arabidopsis carotenoid cleavage dioxygenase family demonstrates the divergent roles of this multifunctional enzyme family. *The Plant Journal* 45:982–93
- Bouvier F, Suires C, Mutterer J, Camara B. 2003. Oxidative remodeling of chromoplast carotenoids: identification of the carotenoid dioxygenase *CsCCD* and *CsZCD* genes involved in Crocus secondary metabolite biogenesis. *The Plant Cell* 15:47–62
- Simkin AJ, Schwartz SH, Auldrige M, Taylor MG, Klee HJ. 2004. The tomato carotenoid cleavage dioxygenase 1 genes contribute to the formation of the flavor volatiles  $\beta$ -ionone, pseudoionone, and geranylacetone. *The Plant Journal* 40:882–92
- Floss DS, Walter MH. 2009. Role of carotenoid cleavage dioxygenase 1 (CCD1) in apocarotenoid biogenesis revisited. *Plant Signaling & Behavior* 4:172–75
- Sun Z, Hans J, Walter MH, Matusova R, Beekwilder J, et al. 2008. Cloning and characterisation of a maize carotenoid cleavage dioxygenase (*ZmCCD1*) and its involvement in the biosynthesis of apocarotenoids with various roles in mutualistic and parasitic interactions. *Planta* 228:789–801
- Yahya M, Berim A, Isaacson T, Marzouk S, Bar E, et al. 2015. Isolation and functional characterization of carotenoid cleavage dioxygenase-1 from *Laurus nobilis* L. (Bay Laurel) fruits. *Journal of Agricultural and Food Chemistry* 63:8275–82
- Huang FC, Horváth G, Molnár P, Turcsi E, Deli J, et al. 2009. Substrate promiscuity of RdCCD1, a carotenoid cleavage oxygenase from *Rosa damascena*. *Phytochemistry* 70:457–64
- Auldrige ME, McCarty DR, Klee HJ. 2006. Plant carotenoid cleavage oxygenases and their apocarotenoid products. *Current Opinion in Plant Biology* 9:315–21
- Varghese R, Kumar SU, Doss CGP, Ramamoorthy S. 2021. Unraveling the versatility of CCD4: metabolic engineering, transcriptomic and computational approaches. *Plant Science* 310:110991

17. Ohmiya A, Kishimoto S, Aida R, Yoshioka S, Sumitomo K. 2006. Carotenoid cleavage dioxygenase (CmCCD4a) contributes to white color formation in chrysanthemum petals. *Plant Physiology* 142:1193–201
18. Watanabe K, Oda-Yamamizo C, Sage-Ono K, Ohmiya A, Ono M. 2018. Alteration of flower colour in *Ipomoea nil* through CRISPR/Cas9-mediated mutagenesis of *carotenoid cleavage dioxygenase 4*. *Transgenic Research* 27:25–38
19. Han Y, Lu M, Yue S, Li K, Dong M, et al. 2022. Comparative methylomics and chromatin accessibility analysis in *Osmanthus fragrans* uncovers regulation of genic transcription and mechanisms of key floral scent production. *Horticulture Research* 9:uhac096
20. Hai NTL, Masuda JI, Miyajima I, Thien NQ, Mojtabehi N, et al. 2012. Involvement of carotenoid cleavage dioxygenase 4 gene in tepal-color change in *Lilium brownii* var. *colchesteri*. *Journal of the Japanese Society for Horticultural Science* 81:366–73
21. Han Y, Wang X, Chen W, Dong M, Yuan W, et al. 2014. Differential expression of carotenoid-related genes determines diversified carotenoid coloration in flower petal of *Osmanthus fragrans*. *Tree Genetics & Genomes* 10:329–38
22. Qian J, Jiang L, Qing H, Chen J, Wan Z, et al. 2022. ZeMYB9 regulates cyanidin synthesis by activating the expression of flavonoid 3'-hydroxylase gene in *Zinnia elegans*. *Frontiers in Plant Science* 13:981086
23. Qian J, Lai W, Jiang L, Zhan H, Zhai M, et al. 2021. Association between differential gene expression and anthocyanin biosynthesis underlying the diverse array of petal colors in *Zinnia elegans*. *Scientia Horticulturae* 277:109809
24. Qing H, Qian J, Chen J, Jiang L, Fu J, et al. 2022. Carotenoid analysis and functional characterization of lycopene cyclases in *Zinnia elegans* L. *Industrial Crops and Products* 188:115724
25. Cunningham FX Jr, Gantt E. 2007. A portfolio of plasmids for identification and analysis of carotenoid pathway enzymes: *Adonis aestivalis* as a case study. *Photosynthesis Research* 92:245–59
26. Lichtenthaler HK. 1987. Chlorophylls and carotenoids: pigments of photosynthetic biomembranes. *Methods in Enzymology* 148:350–82
27. Kumar S, Stecher G, Li M, Knyaz C, Tamura K. 2018. MEGA X: molecular evolutionary genetics analysis across computing platforms. *Molecular Biology and Evolution* 35:1457–549
28. Cunningham FX Jr, Sun Z, Chamovitz D, Hirschberg J, Gantt E. 1994. Molecular structure and enzymatic function of lycopene cyclase from the cyanobacterium *Synechococcus* sp strain PCC7942. *The Plant Cell* 6:1107–21
29. Cunningham FX Jr, Gantt E. 2005. A study in scarlet: enzymes of ketocarotenoid biosynthesis in the flowers of *Adonis aestivalis*. *The Plant Journal* 41:478–92
30. Cunningham FX Jr, Pogson B, Sun Z, McDonald KA, DellaPenna D, et al. 1996. Functional analysis of the beta and epsilon lycopene cyclase enzymes of *Arabidopsis* reveals a mechanism for control of cyclic carotenoid formation. *The Plant Cell* 8:1613–26
31. Livak KJ, Schmittgen TD. 2001. Analysis of relative gene expression data using real-time quantitative PCR and the  $2^{-\Delta\Delta Ct}$  method. *Methods* 25:402–8
32. Ahrazem O, Trapero A, Gómez MD, Rubio-Moraga A, Gómez-Gómez L. 2010. Genomic analysis and gene structure of the plant carotenoid dioxygenase 4 family: a deeper study in *Crocus sativus* and its allies. *Genomics* 96:239–50
33. Kiser PD, Farquhar ER, Shi W, Sui X, Chance MR, et al. 2012. Structure of RPE65 isomerase in a lipidic matrix reveals roles for phospholipids and iron in catalysis. *Proceedings of the National Academy of Sciences of the United States of America* 109:E2747–E2756
34. Kloer DP, Schulz GE. 2006. Structural and biological aspects of carotenoid cleavage. *Cellular and Molecular Life Sciences* 63:2291–303
35. Wang Y, Xu J, Liu A. 2022. Identification of the carotenoid cleavage dioxygenase genes and functional analysis reveal DoCCD1 is potentially involved in beta-ionone formation in *Dendrobium officinale*. *Frontiers in Plant Science* 13:967819
36. Jia K, Baz L, Al-Babili S. 2018. From carotenoids to strigolactones. *Journal of Experimental Botany* 69:2189–204
37. Wang J, Zhang N, Zhao M, Jing T, Jin J, et al. 2020. Carotenoid cleavage dioxygenase 4 catalyzes the formation of carotenoid-derived volatile  $\beta$ -ionone during tea (*Camellia sinensis*) withering. *Journal of Agricultural and Food Chemistry* 68:1684–90
38. Gao J, Yang S, Tang K, Li G, Gao X, et al. 2021. GmCCD4 controls carotenoid content in soybeans. *Plant Biotechnology Journal* 19:801–13
39. Bruno M, Beyer P, Al-Babili S. 2015. The potato carotenoid cleavage dioxygenase 4 catalyzes a single cleavage of  $\beta$ -ionone ring-containing carotenes and non-epoxidated xanthophylls. *Archives of Biochemistry and Biophysics* 572:126–33
40. Rodrigo MJ, Alquézar B, Alós E, Medina V, Carmona L, et al. 2013. A novel carotenoid cleavage activity involved in the biosynthesis of *Citrus* fruit-specific apocarotenoid pigments. *Journal of Experimental Botany* 64:4461–78
41. Ma G, Zhang L, Matsuta A, Matsutani K, Yamawaki K, et al. 2013. Enzymatic formation of  $\beta$ -citraurin from  $\beta$ -cryptoxanthin and zeaxanthin by carotenoid cleavage dioxygenase4 in the flavedo of citrus fruit. *Plant Physiology* 163:682–95
42. Zheng X, Zhu K, Sun Q, Zhang W, Wang X, et al. 2019. Natural variation in CCD4 promoter underpins species-specific evolution of red coloration in citrus peel. *Molecular Plant* 12:1294–307
43. Pacheco SDG, Gasparin AT, Jesus CHA, Sotomaior BB, Ventura ACSB, et al. 2019. Antinociceptive and anti-inflammatory effects of bixin, a carotenoid extracted from the seeds of *Bixa orellana*. *Planta Medica* 85:1216–24
44. Xu Z, Pu X, Gao R, Demurtas OC, Fleck SJ, et al. 2020. Tandem gene duplications drive divergent evolution of caffeine and crocin biosynthetic pathways in plants. *BMC Biology* 18:63
45. He L, Cheng L, Wang J, Liu J, Cheng J, et al. 2022. Carotenoid cleavage dioxygenase 1 catalyzes lutein degradation to influence carotenoid accumulation and color development in foxtail millet grains. *Journal of Agricultural and Food Chemistry* 70:9283–94
46. Ding A, Bao F, Cheng W, Cheng T, Zhang Q. 2023. Phylogeny of PmCCD gene family and expression analysis of flower coloration and stress response in *Prunus mume*. *International Journal of Molecular Sciences* 24:13950
47. Floss DS, Schliemann W, Schmidt J, Strack D, Walter MH. 2008. RNA interference-mediated repression of MtCCD1 in mycorrhizal roots of *Medicago truncatula* causes accumulation of C<sub>27</sub> apocarotenoids, shedding light on the functional role of CCD1. *Plant Physiology* 148:1267–82
48. Bhat ZY, Mohiuddin T, Kumar A, López-Jiménez AJ, Ashraf N. 2021. *Crocus* transcription factors CstMYB1 and CstMYB1R2 modulate apocarotenoid metabolism by regulating carotenogenic genes. *Plant Molecular Biology* 107:49–62
49. Lu C, Qu J, Deng C, Liu F, Zhang F, et al. 2022. The transcription factor complex CmAP3-CmPI-CmUIF1 modulates carotenoid metabolism by directly regulating carotenogenic gene CmCCD4a-2 in chrysanthemum. *Horticulture Research* 19:uhac020
50. Huang H, Gao X, Gao X, Zhang S, Zheng Y, et al. 2022. Flower color mutation, pink to orange, through CmGATA4 - CCD4a-5 module regulates carotenoids degradation in chrysanthemum. *Plant Science* 322:111290
51. Han Y, Wang H, Wang X, Li K, Dong M, et al. 2019. Mechanism of floral scent production in *Osmanthus fragrans* and the production and regulation of its key floral constituents,  $\beta$ -ionone and linalool. *Horticulture Research* 6:106



Copyright: © 2024 by the author(s). Published by Maximum Academic Press, Fayetteville, GA. This article is an open access article distributed under Creative Commons Attribution License (CC BY 4.0), visit <https://creativecommons.org/licenses/by/4.0/>.