

# LC05-136 originates from ROC22, green arising from blue and surpassing blue

## Authors

Qibin Wu, Zhenxiang Li,  
Wenxiang Lu, Fangmei Liang,  
Yuebin Zhang\*, Youxiong Que\*


## Correspondences

[ynzyb@sohu.com](mailto:ynzyb@sohu.com);  
[queyouxiong@126.com](mailto:queyouxiong@126.com)

## In Brief

LC05-136 originates from ROC22, was tailored to suit the specific conditions of Liuzhou city of Guangxi province. It is characterized by high yield, high sugar, highly tolerant to drought and cold stress, strong ratooning ability, and wide adaptability, and is recognized as the flagship of the fifth-generation sugarcane varieties in China. To date, LC05-136, has been accumulatively cultivated across more than 1.67 million hectares, occupying 38% of the sugarcane planting area in Guangxi.

## Graphical abstract



## Highlights

- The agronomic features of LC05-136 are described in detail.
- The breeding strategies and cultivation practices for LC05-136 are dissected.
- This variety won the first prize of Sci. & Technol. Awards issued by Guangxi province.

**Citation:** Wu Q, Li Z, Lu W, Liang F, Zhang Y, et al. 2024. LC05-136 originates from ROC22, green arising from blue and surpassing blue. *Tropical Plants* <https://doi.org/10.48130/tp-0024-0027>

## LC05-136 originates from ROC22, green arising from blue and surpassing blue

Qibin Wu<sup>1</sup>, Zhenxiang Li<sup>1,2</sup>, Wenxiang Lu<sup>3</sup>, Fangmei Liang<sup>3</sup>, Yuebin Zhang<sup>1\*</sup> and Youxiong Que<sup>1,2\*</sup>

<sup>1</sup> National Key Laboratory for Tropical Crop Breeding, Institute of Tropical Bioscience and Biotechnology, Sanya Research Institute, Chinese Academy of Tropical Agricultural Sciences/Sugarcane Research Institute, Yunnan Academy of Agricultural Sciences, Sanya/Kaiyuan 572024/661600, China

<sup>2</sup> Key Laboratory of Sugarcane Biology and Genetic Breeding, Ministry of Agriculture and Rural Affairs, National Engineering Research Center for Sugarcane, College of Agriculture, Fujian Agriculture and Forestry University, Fuzhou 350002, China

<sup>3</sup> Liucheng Sugarcane Research Units, Liuzhou 545000, China

\* Corresponding authors, E-mail: [ynzyb@sohu.com](mailto:ynzyb@sohu.com); [queyouxiong@126.com](mailto:queyouxiong@126.com)

### Abstract

Sugarcane plays a crucial role in the agricultural economy and the livelihood of the people, and the seed industry is the key to maintaining a stable sugar supply. In China, sugarcane varieties have witnessed five rounds of improvement and renewal, which significantly drive the development of sugar industry. Among the fifth-generation sugarcane varieties, LC05-136 was developed by Liucheng Sugarcane Research Units (LC-SRU), with reference to ROC22, and was tailored to suit the specific conditions of the region. This variety, emerged as a flagship, is known for several significant advantages, including high yield, high sugar, highly tolerant to drought and cold stress, strong ratooning ability, and wide adaptability. To date, LC05-136, has been accumulatively cultivated across more than 1.67 million hectares in China. We can reasonably deduce that in sugarcane breeding, selection of "marshals" with a well-balanced combination of desirable traits is far superior to "generals" that excel in only one favorable trait. As it truly deserves, in 2022, the breeding and promotion of LC05-136 won the first prize of Science and Technology Awards issued by the People's Government of Guangxi Zhuang Autonomous Region.

**Citation:** Wu Q, Li Z, Lu W, Liang F, Zhang Y, et al. 2024. LC05-136 originates from ROC22, green arising from blue and surpassing blue. *Tropical Plants* <https://doi.org/10.48130/tp-0024-0027>

Sugarcane, accounting for approximately 90% of total sugar output, is the predominant sugar crop, which plays a crucial role in the agricultural economy and the livelihood of the people in China<sup>[1,2]</sup>. The primary sugarcane cultivation regions include Guangxi, Yunnan, Guangdong, and Hainan provinces. In 2023, sugarcane has been cultivated across roughly 1.11 million hectares, with Guangxi accounting for 0.75 million hectares, Yunnan for 0.26 million hectares, Guangdong for 80,000 hectares, and Hainan for 13,200 hectares. This has resulted in a sugar production of approximately 8.0 million tons. The seed industry, contributing up to 60% to its growth, is a key factor in maintaining a stable sugar supply. Since the establishment of the People's Republic of China, sugarcane varieties have successfully completed five rounds of improvement and renewal, resulting in an average yield increase of approximately 1.0 ton per round and 1% increase in sugar content<sup>[3,4]</sup>. These advancements have significantly driven the continuous, rapid and healthy development of sugar industry in China. In this situation, Liuzhou city of Guangxi province, located in the northernmost sugarcane planting region in China, faces challenges especially poor ecological conditions, including barren dry slopes, cold or frost, and drought. It is generally believed that, if sugarcane can grow well here, it may grow even better elsewhere<sup>[5]</sup>. Recently, Liucheng Sugarcane Research Units (LC-SRU) has developed a new and improved variety named LC05-136 with reference to ROC22, which was tailored to suit the specific conditions of the region (Fig. 1).


In 2005, Wenxiang Lu, director of LC-SRU, developed LC05-136 by selecting from over 100,000 sugarcane seedlings. This

variety, originating from the crossing between CP81-1254 and ROC22<sup>[6]</sup>, is characterized by tall, compact, and moderate growth, featuring medium to large, erect and uniform, and solid stems, cylindrical internodes, straight leaf posture, green leaf color, purple-red leaf sheaths, and easy defoliation<sup>[3,7]</sup>. During the national sugarcane variety regional trial in 2012-2013, LC05-136 exhibited an average yield of 100.74 tons per hectare (t/ha), surpassing the control ROC22 by 0.87%, of which the plant cane increased by 0.15% and the ratoon cane increased by 2.10%. Compared to ROC22, the sugar content averaged at 15.16 t/ha, representing a 5.41% increase (Table 1). Surprisingly, LC05-136 is known for several significant advantages, including but not limited to high yield, high sugar, highly tolerant to drought and cold stress, strong ratooning ability, and wide adaptability (Table 1), making it suitable for cultivation in moderately fertile acreage across major sugarcane planting regions in China, mostly Guangxi, Guangdong, Yunnan, and Hainan provinces<sup>[3,7-9]</sup>. This variety is moderately resistant to sugarcane mosaic disease and sugarcane smut disease in the national sugarcane variety regional trial (2012-2013)<sup>[7,10]</sup>. The recommended row spacing is 100-120 cm, with a general seed amount of 120,000~150,000 buds/ha and the seeds in double or triple buds. On dry slopes, consider planting 3-4 buds in each section. To ensure root retention for over 3 years and enhance planting efficiency, the suitable fertilization time is in late May for plant cane but in mid-April for ratoon cane. As this variety grows rapidly, field management should be timely and thorough. Remember to remove plants infected with smut disease promptly and implement effective measures

**Table 1.** Comprehensive evaluation on variety traits for LC05-136 and ROC22.

Varieties	PH	SD	SSW	ENS	SY	SC	SMDR	SSDR	DT	CT
LC05-136	286.21	2.74	1.64	66573	100.74	15.16	MR	MR	HR	HR
ROC22	300.19	2.72	1.70	64586	99.87	14.39	HS	S	R	MR

The origin data was extracted from Lu and Lu (2015)<sup>[7]</sup>, Luo et al (2016)<sup>[8]</sup> and You et al (2022)<sup>[10]</sup>. PH, plant height (cm); SD, stem diameter (cm); SSW, single stem weight (kg/stem); ENS, effective number of stems (stems/ha); SY, sugarcane yield (t/ha); SC, sugar content (t/ha); SMDR, sugarcane mosaic disease resistance; SSSDR, sugarcane smut disease resistance; DT, drought tolerance; CT, cold tolerance; MR, moderately resistant; HR, highly resistant; HS, highly susceptible. During the national sugarcane variety regional trial in 2012-2013, LC05-136 was moderately resistant to sugarcane smut disease. Unfortunately, the variety has been found to be susceptible to sugarcane smut in recent years.

**Fig. 1** Green LC05-136 originates from blue ROC22 and surpassing blue.

against thrips, aphids, and stem borers<sup>[10]</sup>. Collectively, LC05-136 has been well and widely accepted by sugarcane growers and sugar enterprises (Fig. 1).

Prior to 2013, there is a significant shortage of independently bred elite sugarcane varieties in mainland China<sup>[11]</sup>. ROC22, a variety introduced from Taiwan province, accounted for 55.54% sugarcane planting area nationwide and a striking percentage of 68.20% in Guangxi province<sup>[12]</sup>. In 2014, LC05-136 was successively certified by China Committee for Crop Variety Registration and then by Guangxi Committee for Crop Variety Registration. What excite us all is that, it was greatly promoted, with the rapid expansion of planting area. By 2021, LC05-136 reached 303,333 hectares in mainland China, with Guangxi contributing 278,200 hectares. Fortunately but not surprisingly, it was designated as the popular or mainly promoted sugarcane variety by the Ministry of Agriculture and Rural Development of the People's Republic of China in 2022. To date, LC05-136, emerged as a flagship of the fifth-generation sugarcane varieties, has been accumulatively cultivated across more than 1.67 million hectares in China, occupying 38% of the sugarcane planting area in Guangxi<sup>[4,13,14]</sup>. Despite the success, LC05-136 also exhibits some disadvantages, such as inadequate resistance to lodging and weak ratooning ability in specific regions. Especially, in recent years, the variety has been found to be susceptible to sugarcane smut, with incidence rates increasing with each subsequent ratooning year, opening up a new topic of disease resistance improvement. As the old

saying goes, there is no perfect crop variety, but there are only those that are better. We can reasonably deduce that LC05-136 will eventually be replaced by a new sugarcane variety, reflecting an inevitable progression. This is also the reason why the majority of sugarcane breeders insist on selection and breeding work.

A good sugarcane variety is a collection of traits such as high yield, high sugar, tolerant to abiotic stress, resistant to biotic stress, and strong ratooning ability<sup>[13,15]</sup>. As the cultivation area expands, a minor defect can be magnified indefinitely. Consequently, despite years of breeding efforts, some sugarcane varieties may still greatly suffer from those unfavorable traits such as easy to falling over, vulnerability to insect pests, or low sugar content. Regarding LC05-136, the breeding approach employed has demonstrated the success by utilizing ROC22 as the type of ideal plant and subjecting it to natural local field conditions that well simulate drought and cold stress (Figure 1). In sugarcane breeding, selection of "marshals" with a well-balanced combination of desirable traits is much better than or far superior to "generals" that excel in only one favorable trait<sup>[3]</sup>. Furthermore, involving growers in the selection process well leverages their wealth of experience and ability to quickly identify promising sugarcane varieties. Since one end of sugarcane is connected to growers, while the other is linked to sugar enterprises, the collaborative approach for selection ensures that the needs of growers and enterprises are perfectly combined together and well fulfilled simultaneously. In 2022,

the project "breeding and promotion of sugarcane variety LC05-136" led by Prof. Wenxiang Lu won the first prize of Science and Technology Awards issued by the People's Government of Guangxi Zhuang Autonomous Region. That is the highest confirmation on the acceptance of sugarcane growers, the satisfaction of sugar enterprises, and the increasing efficiency of sugar industry. It truly deserves.

## Author contributions

QW: Formal Analysis, Methodology, Visualization, Funding acquisition, Writing—original draft; ZL: Visualization, Writing—original draft; WL and FL: Project administration, Resources, Writing—review & editing; YZ: Conceptualization, Project administration, Resources, Supervision, Writing—review & editing; YQ: Conceptualization, Funding acquisition, Project administration, Resources, Supervision, Writing—review & editing.

## Data availability

The raw RNA-Seq data of this study were deposited in the Genome Sequence Archive at the China National Center for Bioinformation (<https://www.cncb.ac.cn/>). The original contributions presented in the study are included in the article. Further inquiries can be directed to the corresponding authors.

## Acknowledgments

This work was funded by the Project of National Key Laboratory for Tropical Crop Breeding (NKLTCB-YAAS-2024-S01, Yunnan), Central Public-interest Scientific Institution Basal Research Fund (1630052024003 and 1630052024020), Chinese Academy of Tropical Agricultural Sciences for Science and Technology Innovation Team of National Tropical Agricultural Science Center (CATASCXTD202402), China Agriculture Research System of MOF and MARA (CARS-17), and Yunnan Key Laboratory of Sugarcane Genetic Improvement (2023KFKT001).

## Conflict of interest

The authors declare that they have no conflict of interest.

## Dates

Received 10 May 2024; Accepted 19 June 2024; In press 24 June 2024

## References

1. Chen R, Xu L, Lin Y, Deng Z, Zhang M, Luo J, et al. 2011. *Modern sugarcane genetic breeding*. Beijing: China Agriculture Press, pp: 2–12
2. Lam E, Shine J, Silva JD, Lawton M, Bonos S, Calvino M, et al. 2019. Improving sugarcane for biofuel: engineering for an even better feedstock. *GCB Bioenergy* 1:251–255
3. Zhang Y, Wang L, Lu W, Wu C, Liu J, Zhao P, et al. 2022. *Modern sugarcane breeding theory and variety selection-Heterogeneous complex resistant, high yield and high sugar breeding and application*. Beijing: Science Press
4. Wu Q, Li A, Zhao P, Xia H, Zhang Y, Que Y. 2024. Theory to practice: a success in breeding sugarcane variety YZ08-1609 known as the King of Sugar. *Frontiers in Plant Science* 15:1413108
5. Luo J, Pan Y-B, Que Y, Zhang H, Grisham MP, Xu L. 2015. Biplot evaluation of test environments and identification of mega-environment for sugarcane cultivars in China. *Scientific Reports* 5:15505
6. Chen J, Zhang C, Zhou F, Hu, H, Qiu Y, Wang Q. 2021. Parent analysis of new sugarcane varieties developed in China from 2010 to 2020. *Sugar Crops of China* 43(3):8–12
7. Lu W, Lu L. 2015. Breeding and characteristics of new sugarcane varieties Guiliu 05136. *Sugarcane and Canesugar* 4:1–5
8. Luo Z, Meng Z, He G. 2016. Preliminary report on the evaluation of the cold-tolerant sugarcane variety Liucheng 05136. *Agricultural Technology Service* 5(33):102–103
9. Luo J, Pan Y-B, Xu L, Grisham MP, Zhang H, Que Y. 2015. Rational regional distribution of sugarcane cultivars in China. *Scientific Reports* 5:15721
10. You J, Deng Z, Lu, Y, Wei R, Huang R, Lu W. 2022. Cultivation of high yielding sugarcane Guiliu 05136. *China Seed Industry* 5:129–131
11. Wu Q, Li A, Liu J, Zhao Y, Zhao P, Zhang Y, Que Y. 2024. Sugarcane variety YZ05-51 with high yield and strong resistance: breeding and cultivation perspectives. *Tropical Plants* doi: 10.48130/tp-0024-0019
12. Liang Q, Liu X, Li Y, Lin L, Wang Z, Li X, et al. 2021. Growth and decline of sugarcane cultivar 'ROC22' in Guangxi sugarcane area from 2008 to 2017. *Chinese Journal of Tropical Crops* 42(4):982–990
13. Zhang Y, Zhao P, Hu C, Que Y. 2024. The recent achievements and development trends of sugarcane improvement in China. *China Sugar* 46:87–92
14. Lu G, Liu P, Wu Q, Zhang S, Zhao P, Zhang Y, et al. 2024. Sugarcane breeding: a fantastic past and promising future driven by technology and methods. *Frontiers in Plant Science* 15:1375934
15. Que Y, Wu Q, Zhang H, Luo J, Zhang Y. 2024. Developing new sugarcane varieties suitable for mechanized production in China: principles, strategies and prospects. *Frontiers in Plant Science* 14:1337144



Copyright: © 2024 by the author(s). Published by Maximum Academic Press on behalf of Hainan University. This article is an open access article distributed under Creative Commons Attribution License (CC BY 4.0), visit <https://creativecommons.org/licenses/by/4.0/>.